

# Rough McKean-Vlasov dynamics for robust ensemble Kalman filtering

Michele Coghi<sup>1</sup>, Torstein Nilssen<sup>2</sup>, and Nikolas Nüsken<sup>3</sup>

<sup>1</sup>*Technische Universität Berlin, Straße des 17. Juni 136, 10587 Berlin, Germany, [coghi@tu-berlin.de](mailto:coghi@tu-berlin.de)*

<sup>2</sup>*Institute of Mathematics, University of Agder, 4604 Kristiansand, Norway, [torstein.nilssen@uia.no](mailto:torstein.nilssen@uia.no)*

<sup>3</sup>*Institute of Mathematics, Universität Potsdam, 14476 Potsdam, Germany, [nuesken@uni-potsdam.de](mailto:nuesken@uni-potsdam.de)*

July 15, 2021

## Abstract

Motivated by the challenge of incorporating data into misspecified and multiscale dynamical models, we study a McKean-Vlasov equation that contains the data stream as a common driving rough path. This setting allows us to prove well-posedness as well as continuity with respect to the driver in an appropriate rough-path topology. The latter property is key in our subsequent development of a robust data assimilation methodology: We establish propagation of chaos for the associated interacting particle system, which in turn is suggestive of a numerical scheme that can be viewed as an extension of the ensemble Kalman filter to a rough-path framework. Finally, we discuss a data-driven method based on subsampling to construct suitable rough path lifts and demonstrate the robustness of our scheme in a number of numerical experiments related to parameter estimation problems in multiscale contexts.

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Background in filtering, robust representations and McKean-Vlasov dynamics</b>	<b>5</b>
<b>3</b>	<b>Preliminaries</b>	<b>10</b>
<b>4</b>	<b>Stochastic rough McKean-Vlasov equations</b>	<b>11</b>
<b>5</b>	<b>The interacting particle system</b>	<b>22</b>
<b>6</b>	<b>The numerical scheme, construction of the lift and examples</b>	<b>24</b>
<b>A</b>	<b>From the filter to the McKean-Vlasov</b>	<b>32</b>
<b>B</b>	<b>On the connection between McKean-Vlasov filtering and maximum likelihood estimation</b>	<b>35</b>
<b>C</b>	<b>Bibliography</b>	<b>37</b>

## 1 Introduction

Combining mathematical descriptions of reality with observational data is a key task in economics, science and engineering. In typical applications (such as meteorology [62] or molecular dynamics [70]) there is a hierarchy of models available, ranging from the highly accurate but conceptually and computationally demanding to the approximate but readily interpretable and scalable. Naturally, real-world data is almost always complex, multiscale, and increasingly high-dimensional, whereas the corresponding mathematical abstractions are often preferred to be simple and of comparatively low resolution. The discrepancy between intricate data and reduced-order models poses a significant challenge for their simultaneous treatment, and the failure of some standard statistical approaches in scenarios of this type is well documented [2, 82, 89, 103, 105].

*Robustness to model misspecification and perturbation of the data* is a central concept in the design of statistical methodology capable of bridging scales: Consider a statistical model  $\mathcal{M}_0$  – to be thought of as simple – generating (time-dependent) data  $(Y_t^0)_{t \geq 0}$  and a corresponding algorithmic procedure  $\Phi$  producing the output  $\Phi((Y_t^0)_{t \geq 0})$ . We expect that  $\Phi$  deals adequately with complex data in the case when it is continuous in an appropriate sense. Indeed,  $\mathcal{M}_0$  might be a simplified description of an underlying model family  $(\mathcal{M}_\varepsilon)_{\varepsilon \geq 0}$ , where the formal limit  $\lim_{\varepsilon \rightarrow 0} \mathcal{M}_\varepsilon = \mathcal{M}_0$  encapsulates the passage from a complex to a reduced description. The output  $\Phi((Y_t^\varepsilon)_{t \geq 0})$  on ‘real-world’ data  $(Y_t^\varepsilon)_{t \geq 0}$  is then close to  $\Phi((Y_t^0)_{t \geq 0})$  by continuity even though  $\Phi$  has been constructed on the basis of  $\mathcal{M}_0$ .

Unfortunately, in various contexts (for instance, in parameter estimation for diffusions [37] and stochastic filtering [10]) the map  $\Phi$  is given in terms of stochastic integrals against  $(Y_t^0)_{t \geq 0}$  which are well-known to be discontinuous with respect to standard topologies [48, 50]. The theory of *rough paths* provides a principled route towards constructing continuous modifications  $\Phi'$  of  $\Phi$  (employed, for instance, in [29, 37]) by replacing stochastic integrals in terms of rough integrals defined on appropriately lifted paths  $(Y, \mathbb{Y})$ . Although the difficulty of obtaining or imposing the additional information  $\mathbb{Y}$  is well known (see, however, [9, 44, 71] and Section 6 of this article), recent works have demonstrated the potential of including path signatures into data-driven methods to compress information efficiently or unveil multiscale structure [21, 76].

In this paper, we follow the rough path paradigm just described and develop a robust version of the *Ensemble Kalman Filter* (EnKF) [16, 43], drawing on a reformulation of the Kushner-Stratonovich SPDE for stochastic filtering [10] in terms of McKean-Vlasov dynamics (known as the *feedback particle filter* [84, 96, 100]). The EnKF is a versatile approximate procedure for Bayesian inference that is observed to perform particularly well in high-dimensional settings [92] and has recently been applied to problems in machine learning [54, 90]. Our analysis includes the case when the model and observation noises are correlated as this is precisely the setting in which significant obstacles in the construction of robust filters are known to occur [29]. Furthermore, as shown in [80] and Appendix B, stochastic filtering with correlated noises is a natural generalisation of maximum likelihood parameter estimation for diffusions in the context of inexact measurements. In this context, we discuss the relationship between subsampling-based approaches to multiscale parameter estimation [89] to the task of estimating the Lévy area in a rough paths approach.

The construction put forward in this paper requires the formulation of McKean-Vlasov equations in a rough path setting (as studied first in [19] and more recently in the twin papers [7, 8], both considering the dynamics to be driven by a random rough path). The approach introduced in [27] is more suitable for our needs as it allows a clear separation between independent Brownian motions and a common deterministic noise. However, the assumptions in [27] do not cover unbounded coefficients, while the common noise coefficient may depend on both the state of the solution and its law. In equation (11) considered below, the coefficient  $P$  only depends on the law of  $\hat{X}$ , which simplifies the problem to some extent, but also allows us to use a different approach, where we treat the stochastic and the rough integrals in two separated steps. As it will become clear from the proofs in Section 4.4, there is no need to create a joint rough path (or rough driver). All previous works on rough McKean-Vlasov equations deal with bounded globally Lipschitz-continuous (actually smoother) coefficients. These results cannot be applied here, as the coefficient  $P$  has linear growth in the measure of the solution and is locally-Lipschitz with Lipschitz constant depending on the moments of the solution.

## 1.1 Setting and main results

In this section we specify the exact setting and present our main results. Throughout the paper we fix a filtered probability space  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$  satisfying the usual conditions and consider the following filtering model,

$$dX_t = f(X_t)dt + G^{1/2}dW_t, \quad X_0 \sim \pi_0 \quad (1a)$$

$$dY_t = h(X_t)dt + U dW_t + R^{1/2}dV_t, \quad Y_0 = 0. \quad (1b)$$

In the above display, (1a) represents the (hidden) signal, whereas (1b) specifies the available observations. Note straight away that the signal noise  $G^{1/2}dW_t$  and the observation noise  $U dW_t + R^{1/2}dV_t$  may be correlated. We assume that the signal is  $D$ -dimensional, that is,  $X_t \in \mathbb{R}^D$ , and that the observations are  $d$ -dimensional,  $Y_t \in \mathbb{R}^d$ . In applications it is often the case that  $D \gg d$ . The maps  $f : \mathbb{R}^D \rightarrow \mathbb{R}^D$  and  $h : \mathbb{R}^D \rightarrow \mathbb{R}^d$  are assumed to be sufficiently regular (see below). We assume  $G \in \mathbb{R}^{D \times D}$ ,  $R \in \mathbb{R}^{d \times d}$  to be symmetric nonnegative definite and  $U \in \mathbb{R}^{d \times D}$ . Furthermore,  $(W_t)_{t \geq 0}$  and  $(V_t)_{t \geq 0}$  denote independent  $D$ - and  $d$ -dimensional standard Brownian motions, respectively. Lastly, we assume that the observation covariance

$$C = UU^T + R \in \mathbb{R}^{d \times d} \quad (2)$$

is strictly positive definite. Defining the observation  $\sigma$ -algebras

$$\mathcal{Y}_t = \sigma \{(Y_s)_{0 \leq s \leq t}\} \vee \mathcal{N}, \quad t \geq 0, \quad (3)$$

where  $\mathcal{N} \subset \mathcal{F}$  is the collection of  $\mathbb{P}$ -null sets, our objective is to compute or approximate the filtering measures

$$\pi_t[\phi] = \mathbb{E}[\phi(X_t)|\mathcal{Y}_t], \quad (4)$$

for bounded and measurable test functions  $\phi$ ; we refer to [10] for technical details.

As a step towards tractable numerical approximations of (4), we introduce the following McKean-Vlasov equation,

$$\begin{cases} d\hat{X}_t &= f(\hat{X}_t)dt + G^{1/2}d\hat{W}_t + K_t(\hat{X}_t)C^{-1} \circ dI_t + \Xi_t(\hat{X}_t)dt \\ dI_t &= dY_t - \left(h(\hat{X}_t)dt + U d\hat{W}_t + R^{1/2}d\hat{V}_t\right), \end{cases} \quad (5)$$

where  $(\hat{W}_t)_{t \geq 0}$  (resp.  $(\hat{V}_t)_{t \geq 0}$ ) is a given  $D$ -dimensional (resp.  $d$ -dimensional) Brownian motion, both independent from  $(W_t)_{t \geq 0}$  and  $(V_t)_{t \geq 0}$ . Denoting by  $\hat{\pi} := \mathcal{L}(\hat{X} | \mathcal{Y})$  the conditional law with respect to the common noise  $Y$  of solutions  $\hat{X}$  to (5), the coefficients  $K_t : \mathbb{R}^D \rightarrow \mathbb{R}^{D \times d}$  and  $\Xi_t : \mathbb{R}^D \rightarrow \mathbb{R}^D$  are required to solve the following partial differential equations,

$$\nabla \cdot (\hat{\pi}_t(K_t - BC)) = -\hat{\pi}_t(h - \hat{\pi}_t[h]), \quad (6)$$

and

$$\nabla \cdot (\hat{\pi}_t \Xi_t) = \frac{1}{2} \hat{\pi}_t \left( \text{Trace}(K_t C^{-1} (\nabla h)^T) - \hat{\pi}_t [\text{Trace}(K_t C^{-1} (\nabla h)^T)] \right), \quad (7)$$

for  $t \geq 0$  and  $\mathbb{P}$ -almost surely, where  $B = G^{1/2}U^T C^{-1}$ . Moreover, the common noise  $Y$  coincides with the observation process (1b) and the integral in  $\circ dY$  is understood in the sense of Stratonovich.

The construction of the system (5)-(7) as well as existence of solutions will be detailed in Section 2.4. The filtering problem and the McKean-Vlasov equation (5) are related in the following way.

**Proposition** (formal, see Proposition 2.1). *If  $\pi_t$  admits a density and the solution to (5) is unique, then  $\pi_t = \hat{\pi}_t$ ,  $\mathbb{P}$ -a.s. for every  $t \geq 0$ .*

Reformulations of the filtering problem in terms of McKean-Vlasov dynamics similar to (5)-(7) have been introduced in [66, 100], further analysed in [84], and are commonly referred to as *feedback particle filters*; our formulation combines Stratonovich integration with a stochastic innovation term  $dI_t$  so as to allow for a transition to rough paths in the setting of correlated noises.

One of the practical challenges posed by the system of equations (5)-(7) is solving the PDEs (6) and (7) in  $K$  and  $\Xi$  for a given measure  $\hat{\pi}$ . As has been shown in [96] for a similar system of equations, replacing  $K$  and  $\Xi$  by their best constant-in-space approximations in least-square sense recovers a version of the Ensemble Kalman filter dynamics [16]. We follow this approach (see Lemma 2.2 below) and replace the coefficient  $K_t C^{-1}$  by  $P : \mathcal{P}(\mathbb{R}^D) \rightarrow \mathbb{R}^{D \times d}$ , explicitly defined as

$$P(\pi) := \text{Cov}_\pi(x, h)C^{-1} + B := \pi[x(h - \pi[h])^T]C^{-1} + B, \quad \pi \in \mathcal{P}(\mathbb{R}^D), \quad (8)$$

where  $\mathcal{P}(\mathbb{R}^D)$  refers to the set of probability measures on  $\mathbb{R}^D$ . Similarly, we replace  $\Xi$  by  $\Gamma : \mathcal{P}(\mathbb{R}^D) \rightarrow \mathbb{R}^D$ , defined as

$$\Gamma^\gamma(\pi) = -\frac{1}{2} \text{Trace} \left( P(\pi) \pi[x^\gamma (Dh - \pi[Dh])] \right), \quad 1 \leq \gamma \leq D, \pi \in \mathcal{P}(\mathbb{R}^D), \quad (9)$$

which can be interpreted as an Itô-Stratonovich correction term. Thus, we obtain the system

$$\begin{cases} d\hat{X}_t &= f(\hat{X}_t)dt + G^{1/2}d\hat{W}_t + P(\hat{\pi}_t) \circ dI_t + \Gamma(\hat{\pi}_t)dt \\ dI_t &= dY_t - \left(h(\hat{X}_t)dt + U d\hat{W}_t + R^{1/2}d\hat{V}_t\right). \end{cases} \quad (10)$$

where  $\hat{\pi}_t = \mathcal{L}(\hat{X}_t | \mathcal{Y}_t)$ . This equation is well posed according to Lemma 4.18 below.

In order to construct a robust filter, we replace the common noise in equation (5) by a deterministic rough path  $\mathbf{Y} \in \mathcal{C}^\alpha([0, T], \mathbb{R}^d)$  with regularity  $\frac{1}{3} < \alpha \leq \frac{1}{2}$ . As explained in the introduction, the rationale is that the solutions to rough differential equations are continuous in the rough path driver (see [48]), in contrast to the Itô solution map associated to stochastic differential equations. Moreover, using a deterministic path is natural since we are conditioning on the observation, that we can assume to be given and deterministic.

Applying the modifications from the preceding two paragraphs to equation (5) we obtain the system

$$\begin{cases} d\hat{X}_t &= f(\hat{X}_t)dt + G^{1/2}d\hat{W}_t + P(\hat{\pi}_t) dI_t + \Gamma(\hat{\pi}_t)dt \\ dI_t &= d\mathbf{Y}_t - \left(h(\hat{X}_t)dt + U d\hat{W}_t + R^{1/2}d\hat{V}_t\right). \end{cases} \quad (11)$$

From Lemma 4.13 below, the path  $P(\hat{\pi}_\cdot)$  is controlled by  $\mathbf{Y}$  with Gubinelli derivative [48, Definition 4.6]

$$P(\hat{\pi}_s)' = P(\hat{\pi}_s)^\top \hat{\pi}_s[(x - \pi[x])Dh]^\top C^{-1} = P(\hat{\pi}_s)^\top \text{Cov}_{\hat{\pi}}(x, Dh)C^{-1}, \quad (12)$$

so that the rough integral in equation (11) will make sense (see Section 3.3 below for an overview on controlled rough paths). Moreover, when  $Y$  is a semimartingale with covariance  $\text{Cov}(Y_t, Y_s) = C(t-s)$ , for  $s \leq t$ , the correction between the Stratonovich and Itô rough path lift is given by  $-\frac{1}{2} \text{Trace}(P(\hat{\pi}_t)'C) dt$ , which corresponds to  $\Gamma(\hat{\pi}) dt$ .

Our main result is the following well-posedness and stability theorem for (11).

**Theorem 1.1.** *Let  $1/3 < \alpha < 1/2$  and  $\hat{X}_0 \in L^\rho(\Omega, \mathbb{R}^D)$  with  $\rho > 2/(1-2\alpha)$ . Assume  $h \in C_b^2(\mathbb{R}^D, \mathbb{R}^d)$  and that  $f : \mathbb{R}^D \rightarrow \mathbb{R}^D$  is bounded and Lipschitz-continuous. Then equation (11) admits a unique solution. Moreover, the map  $\mathcal{C}^\alpha([0, T], \mathbb{R}^d) \ni \mathbf{Y} \mapsto \hat{\pi}_t \in \mathcal{P}(C([0, T], \mathbb{R}^D))$  is continuous. Moreover, if  $\mathbf{Y}$  is the Stratonovich lift of  $Y$  in (10), then the solutions to (10) and (11) coincide,  $\mathbb{P}$ -almost surely.*

The proof is a consequence of the results in Section 4.4 and can be found at the end of that section. Associated to the McKean-Vlasov equation (11) we also study the following system of mean-field interacting particles,

$$d\hat{X}_t^i = f(\hat{X}_t^i) dt + G^{1/2} d\hat{W}_t^i + P(\mu_t^N) dI_t^i + \Gamma(\mu_t^N) dt, \quad (13a)$$

$$dI_t^i = d\mathbf{Y}_t - (h(\hat{X}_t^i) dt + U d\hat{W}_t^i + R^{1/2} d\hat{V}_t^i), \quad (13b)$$

where  $\mu_t^N := \frac{1}{N} \sum_{i=1}^N \delta_{X_t^i}$  is the empirical measure of the system and  $\mathbf{Y} \in \mathcal{C}^\alpha([0, T], \mathbb{R}^d)$  is the canonical lift of a differentiable and bounded path  $Y : [0, T] \rightarrow \mathbb{R}^d$  with bounded cadlag derivative. We have the following well-posedness and convergence result for the interacting particles.

**Theorem** (see Remark 5.2 and Theorem 5.8). *Under the same assumptions of Theorem 1.1, let  $(\mathbf{Y}^\delta)_{\delta>0}$  be a family of lifts of bounded differentiable paths with cadlag derivatives that approximate  $\mathbf{Y}$  in the rough-path metric. Let  $\mu^{\delta, N}$  be the empirical measure of (13) driven by  $\mathbf{Y}^\delta$ . Then there exists a sequence  $\delta_N$  such that, for every  $t \in [0, T]$ ,  $\mu_t^{N, \delta_N} \xrightarrow{N \rightarrow \infty} \pi_t$ , in  $\rho$ -Wasserstein distance in  $L^1$ .*

The preceding two theorems suggest that numerical methods based on the interacting particle system (13) are robust to perturbations in the data, and hence suitable for applications in multiscale contexts as described in the introduction. Inspired by Davie's work [31], we propose the following recursive numerical scheme,

$$X_{k+1}^i = X_k^i + f(X_k^i) \Delta t + G^{1/2} \sqrt{\Delta t} \xi_k^i + \hat{P}_k \left( \Delta Y_k - (h(X_k^i) \Delta t + U \sqrt{\Delta t} \xi_k^i + R^{1/2} \sqrt{\Delta t} \eta_k^i) \right) \quad (14a)$$

$$+ \widehat{\text{Cov}}(x, Dh) \hat{P}_k \Delta \mathbb{Y}_k + \hat{\Gamma}_k \Delta t, \quad (14b)$$

in the following referred to as the *Rough-Path Ensemble Kalman Filter* (RP-EnKF). Here,  $\Delta t > 0$  is the step size, and  $(\xi_n^i)$  and  $(\eta_n^i)$  denote independent zero mean Gaussian random variables of dimensions  $D$  and  $d$ , respectively. The precise form of the estimator versions  $\hat{P}_k$ ,  $\widehat{\text{Cov}}$  and  $\hat{\Gamma}_k$  will be detailed in Section 6. Finally, the first term in (14b) is built after the Gubinelli derivative from (12). Crucially, the scheme (14) takes the lifted component  $\Delta \mathbb{Y}_k$  (representing iterated integrals of the path  $(Y_t)_{t \geq 0}$ ) as an input. This dependence allows our methodology to appropriately take into account multiscale structure and other information encoded in the signature, but also necessitates appropriate ways to estimate  $\mathbb{Y}_k$  from data. We will discuss this in more depth in Sections 2.3 and 6, but note here that it is natural to decompose  $\Delta \mathbb{Y}_k$  into its symmetric and skew-symmetric part

$$\Delta \mathbb{Y}_k = \Delta \mathbb{Y}_k^{\text{sym}} + \Delta \mathbb{Y}_k^{\text{skew}}. \quad (15)$$

The form of the symmetric part  $\Delta \mathbb{Y}_k^{\text{sym}}$  is suggested by the requirement that the lifted path is geometric,

$$\Delta \mathbb{Y}_k^{\text{sym}} = \frac{1}{2} (y_{k+1} - y_k) \otimes (y_{k+1} - y_k); \quad (16)$$

in particular this expression can readily be computed from discrete-time observations  $y_k$ . The difficulty thus resides in estimating the Lévy area contributions  $\Delta \mathbb{Y}_k^{\text{skew}}$ . We suggest a subsampling-based method, establishing connections to multiscale parameter estimation as investigated in [89]; see Section 2.3. Other approaches towards obtaining  $\Delta \mathbb{Y}_k$  have been developed in [9, 44, 71]. We would like to stress that although estimating  $\Delta \mathbb{Y}_k^{\text{skew}}$  works reasonably well in our experiments (see Sections 6.1 and 6.2), in many applications it might yield satisfactory results to neglect the skew-symmetric part, that is, to use the approximation  $\Delta \mathbb{Y}_k = \Delta \mathbb{Y}_k^{\text{sym}}$ , either because the observation path is one-dimensional, or because the Lévy area term is comparatively small (see Section 6.3). In these cases, the RP-EnKF can be implemented straightforwardly without additional estimation steps, and represents a robust Stratonovich-version of the EnKF.

**Itô, Stratonovich and rough integrals.** Before concluding this introduction, let us comment on the difference between the RP-EnKF scheme (14) and the more conventional EnKF updates

$$X_{k+1}^i = X_k^i + f(X_k^i) \Delta t + G^{1/2} \sqrt{\Delta t} \xi_k^i + \hat{P}_k \left( \Delta Y_k - (h(X_k^i) \Delta t + U \sqrt{\Delta t} \xi_k^i + R^{1/2} \sqrt{\Delta t} \eta_k^i) \right), \quad (17)$$

see [16, 43, 92]. Clearly, (14) and (17) coincide up to the terms in (14b). This difference can be attributed to alternative perspectives on the underlying continuous-time dynamics. Indeed, (17) corresponds to the Euler-Maruyama discretisation of the McKean-Vlasov SDE

$$d\hat{X}_t = f(\hat{X}_t) dt + G^{1/2} dW_t + P(\hat{\pi}_t) \left( dY_t - \left( h(\hat{X}_t) dt + U d\hat{W}_t + R^{1/2} d\hat{V}_t \right) \right), \quad (18)$$

understood in the sense of Itô. In contrast, the term  $\hat{\Gamma}_k \Delta t$  in (14b) can be seen as an Itô-Stratonovich correction to (18), while the first term in (14b) arises from a Milstein-type approximation scheme for Stratonovich SDEs (or as part of the discrete approximation of rough integrals according to Davies). Importantly, the passage to Stratonovich and ultimately rough integrals is indispensable for the continuity statements in Theorem 1.1 and the numerical robustness of the RP-EnKF scheme demonstrated in Section 6. Our approach of replacing Itô by Stratonovich and subsequently rough integrals mirrors the construction in [37] in the context of maximum likelihood parameter estimation for diffusions. Finally, we would like to emphasise that we performed the Itô-Stratonovich conversion at the level of the exact dynamics (5), and only then apply the constant-in-space approximation in Lemma 2.2.

**Our contributions and structure of the paper.** Our main contributions are as follows:

- We derive the McKean-Vlasov system (5)-(7). Crucially, the dynamics (5)-(7) is given entirely in terms of Stratonovich integrals that allow the construction of robust filtering schemes. Previous works have focussed on either Itô or combined Itô-Stratonovich formulations [80, 91, 96, 100]. While the framework developed in [84] is in terms of Stratonovich integrals, it does not encompass correlated model and observation noise and hence does not extend to our setting of interest.
- We prove well-posedness of the rough McKean-Vlasov dynamics (11) as well as continuity in the rough driver  $\mathbf{Y}$  (see Theorem 1.1).
- We show well-posedness as well as propagation of chaos of the interacting particle approximation (13).
- We suggest the RP-EnKF scheme (14) and in particular devise a subsampling based method to estimate the lift components  $\mathbb{Y}_k$  from data. The robustness of the RP-EnKF (and the nonrobustness of the EnKF) is demonstrated using numerical examples in the context of combined state-parameter estimation.

In Section 2 we review the most relevant results in filtering theory, we motivate the use of the McKean-Vlasov equation and explain the concept of a robust filter. In Section 3 we introduce common notation and recall some background on rough paths. In Section 4.4 we present the analysis of the rough McKean-Vlasov equation (11), which includes well-posedness and stability. In Section 5 we treat the interacting particles system (13) and prove well-posedness and propagation of chaos. In Section 6 we detail the construction of the numerical scheme, including a presentation of our subsampling approaching towards estimating Lévy area. Finally, we conclude the paper with some numerical experiments.

## 2 Background in filtering, robust representations and McKean-Vlasov dynamics

In this section we discuss essential background on robust filtering and put our work into perspective. Section 2.1 will summarise existing approaches towards solving the filtering problem posed by (4), both from a theoretical as well as from an algorithmic perspective. In Section 2.2 we review the challenges to these methods posed by perturbations in the observed data  $(Y_s)_{0 \leq s \leq t}$ , leading to the concept of robustness. In Section 2.3 we draw connections of the McKean-Vlasov approach considered in this paper to maximum likelihood based techniques for stochastic differential equations, in particular to the methods developed in [37]. Finally in Section 2.4 we make our McKean-Vlasov formulation as well as the ensemble Kalman approximation precise.

### 2.1 Solutions to the filtering problem and algorithms

It is well known that the measure  $\pi_t$  defined in (4) is a measurable function of the observation path  $(Y_s)_{0 \leq s \leq t}$  and can be obtained as a solution to the **Kushner-Stratonovich SPDE** [10, Section 3.6],

$$\pi_t[\phi] = \pi_0[\phi] + \int_0^t \pi_s[\mathcal{L}\phi] ds + \int_0^t \left( \pi_s[\phi h^\top] - \pi_s[h^\top] \pi_s[\phi] + \pi_s[(B \nabla \phi)^\top] \right) \left( C^{-1} dY_s - \pi_s[h] ds \right), \quad (19)$$

where

$$\mathcal{L}\phi = f \cdot \nabla \phi + \frac{1}{2} \text{Trace}(G \nabla^2 \phi) \quad (20)$$

denotes the infinitesimal generator associated to the signal process (1a). From the computational viewpoint, numerically solving (19) directly (for instance, using grid-based methods) is usually infeasible, especially when the

dimension  $D$  is large (see [10, Section 8.5] for a discussion). Many algorithmic approaches therefore rely on the simulation of carefully constructed interacting particle systems, positing the corresponding (possibly weighted) empirical measures  $\frac{1}{N} \sum_{i=1}^N \delta_{X_t^i}$  as approximations for  $\pi_t$ .

**Sequential Monte Carlo methods** rely on Bayes' theorem in order to approximate the conditional expectations (4). More precisely, defining the likelihood

$$l_t = \exp \left( \int_0^t h(X_s) \cdot C^{-1} dY_s - \frac{1}{2} \int_0^t h(X_s) \cdot C^{-1} h(X_s) ds \right), \quad (21)$$

the filtering measures admit the representation

$$\pi_t[\phi] = \frac{\mathbb{E}[\phi(X_t) l_t | \mathcal{Y}_t]}{\mathbb{E}[l_t | \mathcal{Y}_t]}, \quad (22)$$

according to the Kallianpur-Striebel formula [10, Proposition 3.16]. Consequently, approximations of  $\pi_t$  can be obtained by sampling from the signal dynamics (1a) in conjunction with appropriate weighting and/or resampling steps on the basis of (21). For detailed accounts, we refer the reader to [40, 36, 92]. While sequential Monte Carlo methods reproduce the filtering measures exactly in the large-particle limit (see, for instance, [10, Theorem 9.15]), they tend to become unstable in high-dimensional settings due to weight collapse: the conditional and unconditional laws of  $X_t$  are often so distinct as to render reweighting-based approaches infeasible due to low effective samples sizes.

**Ensemble Kalman filters (EnKFs)** [92, Section 7.1] can be formulated in terms of interacting or mean-field (McKean-Vlasov) diffusions. In the case when  $U = 0$  (that is, when the signal and observation noises are uncorrelated [93, Section 7.1], [15]) the basic EnKF due to Evensen [17, 42, 43] is given by

$$d\hat{X}_t = f(\hat{X}_t) dt + G^{1/2} d\hat{W}_t + P(\pi_t) C^{-1} \left( dY_t - \left( h(\hat{X}_t) dt + R^{1/2} d\hat{V}_t \right) \right), \quad (23)$$

with

$$P(\pi) = \text{Cov}_\pi(x, h), \quad (24)$$

or a standard particle approximation thereof. The system (23)-(24) is motivated by the fact that the corresponding law reproduces  $\pi_t$  exactly in the linear Gaussian case: If  $\pi_0$  is Gaussian,  $f(x) = Fx$  and  $h(x) = Hx$  for appropriate matrices  $H \in \mathbb{R}^{d \times D}$  and  $F \in \mathbb{R}^{D \times D}$ , then  $\pi_t$  remains Gaussian for all  $t \geq 0$ , and  $\text{Law}(\hat{X}_t) = \pi_t$ . In cases where the preceding conditions are not satisfied, the system (23)-(24) becomes an approximation, the accuracy of which is far from well understood theoretically. However, the ensemble Kalman approach has empirically proven to be both fairly reliable in nonlinear settings as well as scalable to high-dimensional scenarios, and therefore nowadays constitutes one of the workhorses in practical data assimilation tasks [92]. We refer to [16] for a recent review of its theoretical properties.

The recently proposed **feedback particle filters** [100, 96] rely on carefully designed McKean-Vlasov diffusions of the type (23) such that the associated (conditional, nonlinear) Fokker-Planck equation coincides with the Kushner-Stratonovich SPDE (19). By construction, such models are exact, and the conditional laws induced by the solutions to feedback particle filter dynamics provide the filtering measures (4). As an illustration,

$$d\hat{X}_t = f(\hat{X}_t) dt + G^{1/2} d\hat{W}_t + K(\hat{X}_t, \pi_t) C^{-1} \left( dY_t - \left( h(\hat{X}_t) dt + R^{1/2} d\hat{V}_t \right) \right) + \Xi(\hat{X}_t, \pi_t) dt, \quad (25)$$

was suggested in [91] where  $K(\cdot, \pi) = \nabla \phi(\cdot, \pi)$  is determined from the elliptic PDE

$$\nabla \cdot (\pi \nabla \phi) = -\pi(h - \pi[h]), \quad (26)$$

and  $\Xi$  is an appropriate correction term (see Section 2.4 for an in-depth discussion). Clearly, the systems (23)-(24) and (25)-(26) are strongly related in spirit, combining a replication of the signal dynamics (1a) with a data-dependent nudging term so as to match the observations. Reiterating the discussion so far, solutions to (25)-(26) provide exact solutions to the filtering problem (4), while solutions to (23)-(24) lead to approximate ones (except in the linear Gaussian case). However, as  $P$  is given explicitly in (24), the system (23)-(24) lends itself straightforwardly to efficient numerical integration, while the system (25)-(26) poses a formidable numerical challenge in the form of the high-dimensional PDE (26). What is more, well-posedness of systems of the type (25)-(26), with coefficients that depend on the law through the solution of a PDE is currently not well understood. Nevertheless, McKean-Vlasov formulations of the type (25)-(26) conveniently link between the theoretically optimal Kushner-Stratonovich SPDE (19) and the numerically tractable and practically relevant ensemble Kalman dynamics (23)-(24). In this paper, we leverage this viewpoint in order to construct a robust version of (17).

## 2.2 Robust filtering

In order to model and solve real-life problems it is highly desirable that the conditional law  $\pi_t$  (or any numerical approximation thereof) depends continuously on the observation path  $(Y_s)_{0 \leq s \leq t}$ : This property would ensure robustness against misspecification of the underlying signal and observation dynamics (as is typical in reduced-order modeling) as well as against anomalies or artefacts in the collection of the data (such as discretisation errors or perturbation by noise), see [10, Chapter 5] for an overview, [22, 23, 32, 67, 34, 35, 33] and Section 2.3 below. Unfortunately, however, the  $\phi$ -dependent measurable map  $(Y_s(\cdot))_{s \in [0, t]} \mapsto \pi_t[\phi]$  provided by (4) can be shown to be neither unique nor continuous [29] in standard topologies. At a fundamental level, this problem is due to the appearance of stochastic integrals against  $(Y_s)_{0 \leq s \leq t}$  in (19) and (21) which are well known to induce classically discontinuous maps (for instance, in the supremum norm), see [48, 50]. In order to address this issue and to obtain a continuous<sup>1</sup> version of the process  $(\pi_t)_{t \geq 0}$ , Clarke suggested using (stochastic) integration by parts in (21) in order to eliminate the  $dY$ -dependence [22]. Notably, this approach is restricted to the case when the signal and observation noises are independent (that is,  $U = 0$ ), see [22, 23, 67], or the case when the observation is one-dimensional (that is,  $Y_t \in \mathbb{R}$ ), see [32, 34, 35, 33, 45]. Addressing the situation of multidimensional correlated observations, the authors of [29] showed by means of a counter-example (see [29, Example 1]) that continuity as a function of  $(Y_s)_{0 \leq s \leq t}$  is impossible to achieve. Instead, they use rough path lifts  $C([0, T]; \mathbb{R}^d) \ni (Y_s)_{0 \leq s \leq T} \mapsto (\mathbf{Y}_s)_{0 \leq s \leq T} \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^d)$  and establish continuity when  $\pi_t$  is considered as a function of the augmented observation path. Similar ideas have been pursued in [55, Theorem 5.3], putting forward the notion of ‘good’ approximations of the observation path. As the aforementioned works are concerned with the likelihood (21), these lay the foundations for the development of robust sequential Monte Carlo methods as reviewed in Section 2.1, and the recent preprint [30] explores that direction. We would also like to mention the works [38, 57] that allow treating the Zakai SPDE (governing the unnormalised filtering distribution [10, Section 3.5]) in a rough paths framework, however noticing that the numerical treatment of SPDEs is faced with enormous challenges, in particular in high-dimensional settings. Some other works addressing issues in robust or multiscale filtering include [3, 4] (assuming uncertainty in the coefficients) as well as a sequence of works by N. Perkowski and coworkers in the context of averaging and homogenisation [14, 12, 13, 59, 60, 73, 72, 101, 102].

In this paper, we instead construct a robust version of the ensemble Kalman filter (23)-(24) on the basis of its connections to feedback particle formulations as in (25)-(26). Before describing our strategy, we review related work on maximum likelihood parameter estimation for stochastic differential equations.

## 2.3 Parameter estimation and filtering in multiscale systems

In this section we present a prototypical example that illustrates some of the challenges in robust filtering as well as the scope of the methods developed in this paper. Consider the SDE

$$dZ_t = F(Z_t, \theta) dt + dW_t, \quad (27)$$

where  $Z_t \in \mathbb{R}^d$ , and  $F : \mathbb{R}^d \times \Theta \rightarrow \mathbb{R}^d$  is a parameterised drift vector field, with parameter set  $\Theta \subset \mathbb{R}^p$ . The objective is to find the true parameter  $\theta^* \in \Theta$  from a noisy realisation of  $(Z_t)_{t \geq 0}$ , that is, we assume that the observation process is given by

$$dY_t = dZ_t + R^{1/2} dV_t. \quad (28)$$

As before,  $R \in \mathbb{R}^{d \times d}$  denotes the observation noise covariance, and  $(V_t)_{t \geq 0}$  stands for a standard  $d$ -dimensional Brownian motion. The problem setting (27)-(28) can be brought into the form (1) by elevating  $\theta$  to a time-dependent variable, that is, by setting  $X_t = (Z_t, \theta_t) \in \mathbb{R}^{d+p}$ , hence viewing (27)-(28) as a combined state-parameter estimation problem, see [80]. Accordingly,  $f : \mathbb{R}^{d+p} \rightarrow \mathbb{R}^{d+p}$  and  $h : \mathbb{R}^{d+p} \rightarrow \mathbb{R}^d$  are then given as  $f(z, \theta) = (F(z, \theta), 0)$  and  $h(z, \theta) = F(z, \theta)$ , and the matrices  $G \in \mathbb{R}^{(d+p) \times (d+p)}$  and  $U \in \mathbb{R}^{d \times (d+p)}$  take the form

$$G = \begin{pmatrix} I_{d \times d} & 0_{d \times p} \\ 0_{p \times d} & 0_{p \times p} \end{pmatrix}, \quad U = \begin{pmatrix} I_{d \times d} & 0_{p \times p} \end{pmatrix}. \quad (29)$$

Finally, the filtering formulation is completed by specifying a *prior distribution* on the initial condition  $(Z_0, \theta_0)$ . The resulting filtering measures  $\pi_t \in \mathcal{P}(\mathbb{R}^{d+p})$  encode the Bayesian posterior on the combined variable  $(Z_t, \theta_t)$ . Consequently, the  $\theta$ -marginals provide mean a posteriori estimates on the parameter of interest as well as corresponding bounds on Bayesian uncertainty.

In the particular case when the path  $(Z_t)_{t \geq 0}$  is observed without contamination by noise, that is,  $R = 0$ , and  $F$  is linear<sup>2</sup> in  $\theta$ , that is,  $F(z, \theta) = \theta f(z)$  with  $f$  satisfying appropriate nondegeneracy conditions [37], the

<sup>1</sup>As pointed out in [29], the continuity requirement restores the uniqueness of the map  $(Y_s(\cdot))_{s \in [0, t]} \mapsto \pi_t[\phi]$ .

<sup>2</sup>For simplicity of the presentation, we also assume here that  $\theta$  is one-dimensional, that is,  $\Theta \subset \mathbb{R}$ .

parameter  $\theta \in \Theta$  can be recovered from the maximum likelihood estimator

$$\theta_T^*(Z) = \frac{\int_0^T f(Z_t) dZ_t}{\int_0^T |f(Z_t)|^2 dt} \quad (30)$$

in the limit when  $T \rightarrow \infty$ , see [68, 74]. Furthermore, in this case the McKean-Vlasov dynamics suggested in this paper can be solved explicitly, and the corresponding means are directly related to (30), see Appendix B and [80]. It is well known that the estimator (30) can be inaccurate when evaluated on paths that only approximately satisfy (27), for instance when (27) represents a reduced description of an underlying multiscale dynamics [2, 89, 105]. A common approach towards addressing this problem is to subsample the data, see [1, 5, 6, 51, 52, 61, 63, 64, 83, 88] for methodological aspects. For specific applications see [78] (multiscale inverse problems), [2, 82, 105] (economics and finance), and [28, 103] (ocean and atmospheric science).

A different approach towards robustness of the estimator (30) has been taken in [37], addressing the discontinuity of the Itô integral  $\int_0^T f(Z_t) dZ_t$ . To resolve this issue, the authors suggest replacing Itô by Stratonovich integration (motivated by the Wong-Zakai theorem [48, Theorem 9.3] and entailing a correction term involving  $\nabla f$ ), and subsequently using rough paths integration instead of Stratonovich integration (relying on a suitable lift  $C([0, T]; \mathbb{R}^d) \ni (Z_s)_{0 \leq s \leq T} \mapsto (\mathbf{Z}_s)_{0 \leq s \leq T} \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^d)$ ). The resulting estimator

$$\theta_T^{\text{RP}}(\mathbf{Z}) = \frac{\int_0^T f(Z_t) d\mathbf{Z}_t - \frac{1}{2} \int_0^T \text{Trace}(\nabla f)(Z_t) dt}{\int_0^T |f(Z_t)|^2 dt} \quad (31)$$

can then be shown to be continuous as a map from  $\mathcal{C}^\alpha([0, T]; \mathbb{R}^d)$  to  $\mathbb{R}$ . The construction of the RP-EnKF dynamics (14) follows a similar line of reasoning, but our approach is applicable to situations where the path  $(Z_t)_{t \geq 0}$  is contaminated by noise ( $R \neq 0$ ) and where  $F(z, \theta)$  is nonlinear in  $\theta$ . The latter generalisation makes our method suitable to applications involving deep learning, that is, when the drift in (27) is parameterised by a neural network as in [54]. We discuss the idea of subsampling the observed data path in the context of constructing an appropriate rough path lift in Remark 6.2 below in Section 6.

## 2.4 From the filtering problem to the McKean-Vlasov equation

In this section, we discuss the construction of the McKean-Vlasov system (5)-(7) and the nature of the approximation in (10) and (11). The general idea goes back to [66, 100], and various modifications have been proposed in [80, 84, 91]. Our formulation combines Stratonovich integration (as in [84]) in order to later invoke Wong-Zakai type approximation results with a stochastic innovation term (as in [80, 92]) as required for the case of correlated model and observation noise. For the sake of clarity, we repeat the PDEs (6) and (7) in their respective index forms (using Einstein's summation convention),

$$\partial_i \left( \hat{\pi}_t \left( K_t^{ij} - (BC)^{ij} \right) \right) = -\hat{\pi}_t \left( h^j - \hat{\pi}_t[h^j] \right), \quad j = 1, \dots, d, \quad (32)$$

and

$$\partial_i (\hat{\pi}_t \Xi_t^i) = \frac{1}{2} \hat{\pi}_t \left( (K_t C^{-1})^{ij} \partial_i h^j - \hat{\pi}_t[(K_t C^{-1})^{ij} \partial_i h^j] \right). \quad (33)$$

The McKean-Vlasov system (5)-(7) solves the filtering problem in the following sense:

**Proposition 2.1.** *Let  $T > 0$ , assume that the system (1) admits a unique solution  $(X_t, Y_t)$  and that the Zakai equation associated to the corresponding filtering problem is well posed. Let  $\pi$  be the conditional law of  $X$  given  $Y$  as defined in (4) and assume that  $\pi_t$  admits a  $C^1$ -density with respect to the Lebesgue measure,  $\mathbb{P}$ -a.s., for all  $t \in [0, T]$ . Moreover, assume that the McKean-Vlasov equation (5) admits a unique solution  $(\hat{X}_t)_{t \in [0, T]}$  such that its conditional law  $\hat{\pi}_t$  admits a  $C^1$ -density with respect to the Lebesgue measure,  $\mathbb{P}$ -a.s.. Assume that  $K$  and  $\Xi$  are predictable stochastic processes with values in  $C^1(\mathbb{R}^D; \mathbb{R}^{D \times d})$  and  $C^1(\mathbb{R}^D; \mathbb{R}^D)$ , respectively, independent from  $(\hat{V}_t)_{t \geq 0}$  and  $(\hat{W}_t)_{t \geq 0}$ , and such that the PDEs (6) and (7) are satisfied,  $\mathbb{P}$ -a.s.. Then  $\pi_t = \hat{\pi}_t$ , for all  $t \in [0, T]$ .*

To prove Proposition 2.1, we define the unnormalised conditional law associated to the McKean-Vlasov dynamics (5),

$$\hat{\rho}_t[\phi] := \mathbb{E}[\phi(X_t) l_t | \mathcal{Y}_t], \quad (34)$$

where the likelihood  $l_t$  is defined in (21). Comparing the evolution of  $\hat{\rho}_t$  with the solution of the Zakai equation [10, Section 3.5] allows us to derive the PDEs (6) and (7). This approach allows us to circumvent the stringent regularity condition in [84, Assumption 3.4]. For details see Appendix A.

One of the numerical challenges posed by the system of equations (5)-(7) is to obtain (approximate) solutions  $K$  and  $\Xi$  to the PDEs (6) and (7). We sidestep this problem by using constant-in-space approximations leading to the system (10) of ensemble Kalman filter type. A similar correspondence has been observed in [96] and is optimal in the following sense:

**Lemma 2.2.** Let  $\pi \in \mathcal{P}(\mathbb{R}^D)$  and  $K : \mathbb{R}^D \rightarrow \mathbb{R}^{D \times d}$  be a (weak) solution to (6). Denote by  $\tilde{K}$  the best constant approximation in least-squares sense, that is

$$\tilde{K} = \arg \min_{\tilde{K} \in \mathbb{R}^{D \times d}} \int_{\mathbb{R}^D} \|K(x) - \tilde{K}\|_F^2 d\pi(x), \quad (35)$$

where  $\|\cdot\|_F$  denotes the Frobenius norm. Then  $\tilde{K}$  is given by

$$\tilde{K} = \int_{\mathbb{R}^D} x (h(x) - \pi[h])^\top d\pi(x) - BC \in \mathbb{R}^{D \times d}. \quad (36)$$

Moreover, let  $\Xi$  be a solution to (7) with  $K$  replaced by  $\tilde{K}$ . Then the best constant approximation of  $\Xi$  in least-squares sense (with respect to the Euclidean norm) is given by

$$\tilde{\Xi}^\gamma = -\frac{1}{2} \text{Trace}(\tilde{K} C^{-1} \pi[x^\gamma (Dh - \pi[Dh])]), \quad 1 \leq \gamma \leq D.$$

*Proof.* The variance (35) is minimised when  $\tilde{K} = \pi[K]$ . To compute the expectation of  $K$ , we multiply equation (6) on both sides by  $x \in \mathbb{R}^D$  and integrate by parts to obtain

$$\pi[K] - BC = \int_{\mathbb{R}^D} (K(x) - BC) d\pi(x) = \int_{\mathbb{R}^D} x (h(x) - \pi[h])^\top d\pi(x).$$

Hence,  $\tilde{K} = \pi[K] = \text{Cov}_\pi(x, h) - BC$ .

We now show the corresponding statement for  $\tilde{\Xi}$  when  $K$  is replaced by  $\tilde{K}$  in (7). Again  $\hat{\Xi} = \mathbb{E}[\Xi]$ , since the expectation minimises the variance. For  $1 \leq \gamma \leq D$ , we input the test function  $\phi(x) = x^\gamma$  into the weak formulation (117) of (7) to obtain the desired expression for  $\tilde{\Xi}$ .  $\square$

Constructing numerical approximations for (6)-(7) beyond the constant-in-space approximation is a topic of ongoing research. We mention in particular the approach developed in [97] and analysed in [85] based on diffusion maps as well as the method from [79] based on the Stein geometry [41, 81].

## 2.5 Literature on McKean-Vlasov dynamics

McKean-Vlasov equations are stochastic differential equations whose coefficients depend on the law of the solution. They are sometimes called law-dependent equations. McKean-Vlasov equations have been the subject of several studies starting from the seminal work of McKean [77] and Dobrushin [39]. McKean-Vlasov equations arise as limit of mean-field interacting particle systems, when the number of particles goes to infinity. For a general introduction on the topic we refer the reader to Sznitman [95].

In recent years there has been an increased interest in mean-field particles with common noise, see [25, 65, 66] or [18] in the case of mean-field games. In these type of systems the particles are subject to the same random perturbation and possibly additional independent noises. There is no averaging effect of the common perturbation when the number of particles increases. The limit object is again a law-dependent SDE, but this time the coefficient depends on the conditional law of the solution given the common noise. This is the case for equation (11), where the coefficients  $P$  and  $\Gamma$  depend on the law of  $X$  given  $Y$ .

McKean-Vlasov equations from a rough path perspective were studied for the first time in [19] and more recently in the twin papers [7, 8]. In both of these works the equation is driven by a random rough path that is quite general and can describe the independent noise, the common noise or both. This gives the additional difficulty of needing to keep track of the rough path as a  $L^p$ -valued path. In [19] only the drift of the equation depends on the law of the solution, the coefficients in front of the noise depend only on the state of the solution. The more recent work [8] generalises that approach to include law-dependent coefficients. The authors use the approach by Gubinelli on controlled rough paths (see Section 3.2 for a brief introduction on the topic). In order to do this, they need Lions' approach to calculus in measure spaces endowed with the Wasserstein metric. The equation is then solved as a fixed-point in the mixed  $\mathbb{R}^d$  and  $L^p$ -space. In [24] the case of pathwise McKean-Vlasov equation with additive noise is considered. The basic techniques used are similar to the ones used in the rough-path case, but the need for rough paths is removed thanks to the additive noise. See also [98].

McKean-Vlasov equations with a rough common noise have been studied recently in [27]. This is the first time that a rough McKean-Vlasov equation is studied when there is a clear separation between independent Brownian motions and a common deterministic noise. In [27], the common noise coefficient depends on both the state of the solution and its law. In equation (11), the coefficient  $P$  only depends on the law of  $\hat{X}$ , which simplifies the problem to some extent, but also allows us to use a different approach, where we treat the stochastic and the rough integrals in two separated steps. As it will be clear from the proofs in Section 4.4, there is no need to create a joint rough path (or rough driver). All previous works on rough McKean-Vlasov equations deal with bounded coefficients, which cannot be applied here, as the coefficient  $P$  has linear growth in the measure of the solution.

Very recently the authors of [49] developed a theory of mixed rough and stochastic differential equations under Lipschitz and boundedness conditions on the coefficients and they plan to address the application to McKean-Vlasov equations with common noise in a forthcoming paper.

### 3 Preliminaries

#### 3.1 Notation

Given a metric space  $(S, d_S)$ , we call  $\mathcal{P}(S)$  the space of probability measures on  $S$ . For  $\pi \in \mathcal{P}(S)$  we denote by  $\pi[x] = \int_S x \pi(dx)$  the mean of  $\pi$  and by  $\pi[\phi] = \int_S \phi(x) \pi(dx)$  the integral of a measurable function  $\phi : S \rightarrow \mathbb{R}$  in  $\pi$ .

Let  $\rho > 0$ , if  $S$  is a normed space with norm  $|\cdot|$  and  $\pi \in \mathcal{P}(S)$ , we denote by

$$M^\rho(\pi) = \int_S |x|^\rho \pi(dx), \quad \overline{M}^\rho(\pi) = \int_S |x - \pi[x]|^\rho \pi(dx),$$

the  $\rho$ -moment of  $\pi$  and the  $\rho$ -central moment of  $\pi$ , respectively. For  $\rho \geq 1$ , we call  $\mathcal{P}_\rho(S) \subset \mathcal{P}(S)$  the space of probability measures  $\pi$  on  $S$  such that  $M^\rho(\pi) < \infty$ . We endow this space with the  $\rho$ -Wasserstein metric

$$W_{\rho,S}^\rho(\mu, \nu) := \inf_{m \in \Gamma(\mu, \nu)} \iint_{S \times S} d_S^\rho(x, y) m(dx, dy),$$

where  $\Gamma(\mu, \nu)$  is the set couplings between  $\mu$  and  $\nu$ . For ease of notation, we denote by  $W_\rho$  the Wasserstein metric on  $\mathbb{R}^D$  and by  $W_{\rho,[0,T]}$  the Wasserstein metric on  $(C([0, T], \mathbb{R}^D), \|\cdot\|_\infty)$ . Given a function  $\varphi \in C(\mathbb{R}^m, \mathbb{R}^n)$  and a path  $x \in C([0, T], \mathbb{R}^m)$ , we define

$$[\varphi]_{s,t}^{k,x} := \int_0^1 (1-\theta)^{k-1} \varphi(x_s + \theta \delta x_{s,t}) d\theta,$$

where  $\delta x_{s,t} := x_t - x_s$ . If  $\varphi \in C^1(\mathbb{R}^m, \mathbb{R}^n)$ , we use the following notation for the Taylor expansion

$$\delta \varphi(x)_{s,t} = [D\varphi]_{s,t}^{1,x} \delta x_{s,t}, \quad [\varphi]_{s,t}^{1,x} - \varphi(x_s) = [D\varphi]_{s,t}^{2,x} \delta x_{s,t}.$$

Throughout the paper we use  $D\varphi$  for the usual Fréchet derivative and  $\nabla \varphi = D\varphi^\top$ . We sometimes use the notation  $L_\omega^\rho := L^\rho(\Omega, \mathbb{R}^d)$ . For a stochastic process  $X$  on a filtered probability space  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ , we denote the conditional expectation by  $\mathbb{E}_s[X_t] := \mathbb{E}[X_t | \mathcal{F}_s]$ .

#### 3.2 Background in rough paths

The theory of rough paths is a framework that allows well-posedness and stability properties for equations of the form

$$\dot{X}_t = \xi(X_t) \dot{Y}_t, \quad X_0 \in \mathbb{R}^d, \quad (37)$$

where  $Y$  is a path of regularity lower than the regularity assumptions amenable to classical calculus. For  $\alpha > 0$ , we denote by  $C_2^\alpha([0, T]; \mathbb{R}^d)$  the set of all continuous functions

$$g : \{(s, t) \in [0, T]^2 : s < t\} \rightarrow \mathbb{R}^d$$

such that there exists a constant  $C$  with  $|g_{s,t}| \leq C|t-s|^\alpha$  and we denote by  $[g]_\alpha$  the infimum over all such constants. We denote by  $\|g\|_\alpha := [g]_\alpha + |g_0|$  the  $\alpha$ -Hölder norm. We write  $C^\alpha([0, T]; \mathbb{R}^d)$  for the set of paths  $f : [0, T] \rightarrow \mathbb{R}^d$  such that  $\delta f \in C_2^\alpha([0, T]; \mathbb{R}^d)$ , where we have defined  $\delta f_{s,t} := f_t - f_s$ .

A *rough path* is a pair  $\mathbf{Y} = (Y, \mathbb{Y}) \in \mathcal{C}^\alpha([0, T]; \mathbb{R}^d) \subset C^\alpha([0, T]; \mathbb{R}^d) \times C_2^{2\alpha}([0, T]; \mathbb{R}^{d \times d})$  such that

$$\mathbb{Y}_{s,t} - \mathbb{Y}_{s,u} - \mathbb{Y}_{u,t} = Y_{s,u} \otimes Y_{u,t}. \quad (38)$$

We equip  $\mathcal{C}^\alpha([0, T]; \mathbb{R}^d)$  with its subset topology which we shall call the rough path topology. Relation (38), commonly referred to as Chen's relation, encodes the algebraic property between a path and its *iterated integral*, viz the formal equality

$$\mathbb{Y}_{s,t} = \int_s^t Y_{s,r} \otimes dY_r.$$

When  $\alpha \in (\frac{1}{3}, \frac{1}{2}]$  the above integral is in general not canonically defined using functional analysis. However, in the case of  $Y$  being a sample path of the Brownian motion,  $B$ , one can use probability theory to define iterated integrals using e.g. Itô integration or Stratonovich integration. We denote by  $\mathbf{B} := (B, \mathbb{B}^{Itô}) := (B, \int B \otimes dB)$  and  $\mathbf{B}^{Strat} := (B, \mathbb{B}^{Strat}) := (B, \int B \otimes \circ dB)$  these (random) rough paths, respectively. It is classical that

$$\mathbb{B}_{s,t}^{Itô} = \mathbb{B}_{s,t}^{Strat} + \frac{1}{2}(t-s)I_{d \times d}. \quad (39)$$

The Stratonovich rough path is an example of a *geometric* rough path, that is to say it is in the closure in the rough path topology of the image of the mapping

$$Y \mapsto (Y, \int Y \otimes dY)$$

defined on  $BV([0, T]; \mathbb{R}^d)$ .

Given two rough paths  $\mathbf{Y}^1, \mathbf{Y}^2 \in \mathcal{C}^\alpha([0, T], \mathbb{R}^d)$  we define the following distance,

$$\rho_\alpha(\mathbf{Y}^1, \mathbf{Y}^2) := \|Y^1 - Y^2\|_\alpha + \|\mathbb{Y}^1 - \mathbb{Y}^2\|_{2\alpha}.$$

We refer the reader to [48] for a more comprehensive discussion of the rough paths notations and concepts used in this paper.

### 3.3 Controlled rough paths and rough differential equations

To use rough paths for a solution theory of equations of the form (37), which we rewrite with the formal expression

$$dX_t = \xi(X_t) d\mathbf{Y}_t, \quad (40)$$

we start with the ansatz that the solution  $X$  takes the form of a Taylor-like expansion

$$\delta X_{s,t} = X'_s \delta Y_{s,t} + X^\sharp_{s,t}, \quad (41)$$

where  $X^\sharp$  is of higher regularity than  $X$ , and  $X'$  is the so-called Gubinelli derivative. We denote by  $\mathcal{D}_Y^{2\alpha}([0, T]; \mathbb{R}^n)$  the set of all pairs  $(X, X')$  such that  $X^\sharp$  implicitly defined via (41) satisfies  $(X', X^\sharp) \in C^\alpha([0, T]; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^n)) \times C^{2\alpha}_2([0, T]; \mathbb{R}^n)$ , which also induces the topology on  $\mathcal{D}_Y^{2\alpha}([0, T]; \mathbb{R}^n)$ .

The sewing lemma provides a continuous integration mapping

$$\begin{aligned} \mathcal{D}_Y^{2\alpha}([0, T]; \mathcal{L}(\mathbb{R}^n, \mathbb{R}^d)) &\longrightarrow \mathcal{D}_Y^{2\alpha}([0, T]; \mathbb{R}^n) \\ (X, X') &\longmapsto \left( \int X_r d\mathbf{Y}_r, X \right), \end{aligned}$$

where

$$X^\natural_{s,t} := \int_s^t X_r d\mathbf{Y}_r - X_s Y_{s,t} - X'_s \mathbb{Y}_{s,t}$$

satisfies  $|X^\natural_{s,t}| \leq C|t - s|^{3\alpha}$  for some constant  $C$  only depending on  $\alpha$ . A solution of (40) can now be defined as a fixed point of the composition of the mappings

$$\begin{aligned} \mathcal{D}_Y^{2\alpha}([0, T]; \mathbb{R}^n) &\longrightarrow \mathcal{D}_Y^{2\alpha}([0, T]; \mathcal{L}(\mathbb{R}^n, \mathbb{R}^d)) &\longrightarrow \mathcal{D}_Y^{2\alpha}([0, T]; \mathbb{R}^n) \\ (X, X') &\longmapsto (\xi(X), \xi(X')) = (\xi(X), \nabla \xi(X) X') &\longmapsto \left( \int \xi(X_r) d\mathbf{Y}_r, \xi(X) \right). \end{aligned}$$

From the sewing lemma and the definition of the integration mapping we see that we could equivalently define the solution of (40) as a path  $X : [0, T] \rightarrow \mathbb{R}^n$  such that

$$X^\natural_{s,t} := \delta X_{s,t} - \xi(X_s) Y_{s,t} - \nabla \xi(X_s) \xi(X_s) \mathbb{Y}_{s,t}$$

satisfies  $|X^\natural_{s,t}| \leq C|t - s|^{3\alpha}$ . The latter formulation is usually referred to as Davie's expansion/solution.

One of the remarkable properties of rough path equations is the continuity of the Itô-Lyons map,

$$\begin{aligned} \mathcal{C}_Y^\alpha([0, T]; \mathbb{R}^d) &\longrightarrow C^\alpha([0, T]; \mathbb{R}^n) \\ \mathbf{Y} &\longmapsto X \end{aligned}$$

where  $X = X^\mathbf{Y}$  denotes the solution of (40), provided  $f$  is regular enough. In fact, Theorem 1.1 is an analogous result for the McKean-Vlasov dynamics of the ensemble Kalman filter treated in this paper.

## 4 Stochastic rough McKean-Vlasov equations

In this section, unless otherwise specified we fix  $T > 0$  and  $\rho \geq 1$ . Moreover,  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$  is a complete filtered probability space that supports a standard  $m$ -dimensional Brownian motion  $W$ . Recall that for  $\pi \in \mathcal{P}_\rho(\mathbb{R}^D)$ , we call  $M^\rho(\pi)$  the  $\rho$ -moment of  $\pi$  and  $\bar{M}^\rho(\pi)$  the  $\rho$ -central moment of  $\pi$ .

#### 4.1 McKean-Vlasov SDEs with linear growth in the measure

Consider the measurable functions  $b : [0, T] \times \mathbb{R}^D \times \mathcal{P}_\rho(\mathbb{R}^D) \rightarrow \mathbb{R}^D$  and  $\sigma : [0, T] \times \mathbb{R}^D \times \mathcal{P}_\rho(\mathbb{R}^D) \rightarrow \mathbb{R}^{D \times m}$  satisfying the following assumptions:

**Assumption 1.** Assume that there exists a constant  $C > 0$  such that

(i) (linear growth)  $\forall t \in [0, T], \forall x \in \mathbb{R}^D, \forall \pi \in \mathcal{P}_\rho(\mathbb{R}^D)$ ,

$$|b(t, x, \pi)|, |\sigma(t, x, \pi)| \leq C(1 + M^\rho(\pi))^{\frac{1}{\rho}}, \quad (42)$$

(ii) (locally Lipschitz)  $\forall t \in [0, T], \forall x, y \in \mathbb{R}^D, \forall \pi, \nu \in \mathcal{P}_\rho(\mathbb{R}^D)$ ,

$$|b(t, x, \pi) - b(t, y, \nu)|, |\sigma(t, x, \pi) - \sigma(t, y, \nu)| \leq C(1 + M^\rho(\pi))^{\frac{1}{\rho}} (|x - y| + W_{\rho, \mathbb{R}^D}(\pi, \nu)). \quad (43)$$

*Remark 4.1.* Notice that, if  $\rho \geq \rho'$ , Assumption 1 with  $\rho'$  implies Assumption 1 with  $\rho$ .

Consider the following McKean-Vlasov equation,

$$dX_t = b(t, X_t, \mathcal{L}(X_t)) dt + \sigma(t, X_t, \mathcal{L}(X_t)) dW_t, \quad X_t|_{t=0} = X_0 \in L_\omega^2, \quad (44)$$

where  $W$  is an  $m$ -dimensional Brownian motion.

**Lemma 4.2.** Under Assumption 1 with  $\rho \geq 2$ , equation (44) admits a unique strong solution in the classical sense. Moreover, the solution has continuous sample paths.

*Proof.* To prove well-posedness we use a Picard-type argument. For  $T \geq 0$  and  $K > 2M^2(\pi_0) \vee 1$ , we define

$$B_K := \left\{ \pi \in \mathcal{P}(C([0, T], \mathbb{R}^D)) \mid M^2(\pi_t) \leq K \forall t \in [0, T], M^2(\pi_0) \leq K/2 \right\}.$$

Let  $\pi \in B_K$ . We define the following stochastic differential equation

$$dX_t = b(t, X_t, \pi_t) dt + \sigma(t, X_t, \pi_t) dW_t, \quad \tilde{X}_t|_{t=0} = X_0 \sim \pi_0. \quad (45)$$

By our choice of  $\pi$  and Assumption 1, the coefficients  $b$  and  $\sigma$  are bounded and Lipschitz. By standard theory this equation admits a unique strong solution on the interval  $[0, T]$ , which we denote by  $X^\pi$ . This solution has continuous sample paths,  $\mathbb{P}$ -a.s.. We are ready to define the map

$$\begin{aligned} \Phi : B_K &\rightarrow \mathcal{P}(C([0, T], \mathbb{R}^D)) \\ \pi &\mapsto \mathcal{L}(X^\pi). \end{aligned}$$

Using standard stochastic calculus estimates, Assumption 1 and Gronwall's lemma we obtain the following upper bound on the second moment,

$$\left\| \sup_{t \in [0, T]} |X_t| \right\|_{L_\omega^2} \leq K/2 + TC(1 + K), \quad (46)$$

where  $C > 0$  is a generic constant that only depends on the coefficients  $b$  and  $\sigma$ . Let us now take  $T = 1/(4C)$ , with this choice we have  $\left\| \sup_{t \in [0, T]} |X_t| \right\|_{L_\omega^2} \leq K$ . This implies  $\Phi(B_K) \subset B_K$ . Using again stochastic calculus and Gronwall's lemma, we have that for each  $t \in [0, T]$ ,

$$W_{2, C([0, t], \mathbb{R}^D)}(\Phi(\pi), \Phi(\pi'))^2 \leq \left\| \sup_{s \in [0, t]} |X_s^\pi - X_s^{\pi'}| \right\|_{L_\omega^2}^2 \leq C(1 + K)e^{C(1+K)t} \int_0^t W_{2, \mathbb{R}^D}(\pi_s, \pi'_s)^2 ds, \quad \pi, \pi' \in B_K. \quad (47)$$

Here  $C > 0$  is again a generic constant depending only on  $b$  and  $\sigma$ , possibly different than before. Also notice that  $\pi, \pi'$  are measures on the path space  $C([0, T], \mathbb{R}^D)$  and in the left-hand side of (47) we are considering, with an abuse of notations, their projections on  $C([0, t], \mathbb{R}^D)$ .

Iterating  $n$  times the inequality (47), we obtain

$$W_{2, C([0, T], \mathbb{R}^D)}(\Phi(\pi), \Phi(\pi')) \leq \frac{e^{(n+1)C(1+K)T}}{n!} W_{2, C([0, T], \mathbb{R}^D)}(\pi, \pi'), \quad \pi, \pi' \in B_K.$$

For the choice of  $T = 1/(4C)$ , if we take  $n$  large enough, we have that the map  $\Phi$  is a contraction on  $B_K$ . By the Banach fixed point theorem,  $\Phi$  has a unique fixed point on  $B_K$ , which is the unique solution to equation (44) up to time  $T$ .

Global existence and uniqueness follow by iterating this argument on intervals of fixed length  $1/(4C)$ , which does not depend on the value of the initial condition, but only on the assumptions on the coefficients of the equation.  $\square$

## 4.2 The common noise case

Consider measurable functions  $b$  and  $\sigma$  as in the previous section and  $\beta : [0, T] \times \mathcal{P}_\rho(\mathbb{R}^D) \rightarrow \mathbb{R}^{D \times m_1}$ , each of which satisfying the following assumption on their respective domain.

**Assumption 2.** Let  $(V, |\cdot|)$  be a Banach space, for  $f : [0, T] \times \mathbb{R}^D \times \mathcal{P}_\rho(\mathbb{R}^D) \rightarrow V$  assume that there exists a constant  $C > 0$  such that

(i) (linear growth)  $\forall x \in \mathbb{R}^D, \forall \pi \in \mathcal{P}_\rho(\mathbb{R}^D)$ ,

$$|f(t, x, \pi)| \leq C(1 + \overline{M}^\rho(\pi))^{\frac{1}{\rho}}, \quad (48)$$

(ii) (locally Lipschitz)  $\forall x, y \in \mathbb{R}^D, \forall \pi, \nu \in \mathcal{P}_\rho(\mathbb{R}^D)$ ,

$$|f(t, x, \pi) - f(t, y, \nu)| \leq C(1 + \overline{M}^\rho(\pi))^{\frac{1}{\rho}} (|x - y| + W_{\rho, \mathbb{R}^D}(\pi, \nu)). \quad (49)$$

Notice that, since  $\beta$  is independent of the space variable  $x \in \mathbb{R}^D$ , condition 2 (ii) reduces to local Lipschitz continuity in the measure variable.

Consider the following McKean-Vlasov equation,

$$dX_t = b(t, X_t, \mathcal{L}(X_t | \mathcal{B}_t)) dt + \sigma(t, X_t, \mathcal{L}(X_t | \mathcal{B}_t)) dW_t + \beta(t, \mathcal{L}(X_t | \mathcal{B}_t)) dB_t, \quad X_t|_{t=0} = X_0 \in L_\omega^2, \quad (50)$$

where  $W$  is the  $m$ -dimensional Brownian motion fixed at the beginning of the section and  $B$  is an  $m_1$ -dimensional Brownian motion adapted to  $(\mathcal{F}_t)_{t \geq 0}$ . Assume that  $X_0, W, B$  are independent. In (50),  $\mathcal{L}(X | \mathcal{B}_t)$  is the conditional law of the solution  $X$  given the filtration  $\mathcal{B}_t := \sigma(B_s | 0 \leq s \leq t)$  generated by the common noise  $B$ .

**Lemma 4.3.** Assume Assumption 2. Let  $X$  and  $Z$  be any two solutions to equation (50). Then  $X$  and  $Z$  are indistinguishable.

*Proof.* It follows from the independence of  $W, B$  and  $X_0$  that

$$\mathbb{E}[X_t | \mathcal{B}_t] = \mathbb{E}[X_0] + \int_0^t \mathbb{E}[b(s, X_s, \mathcal{L}(X_s | \mathcal{B}_s)) | \mathcal{B}_s] ds + \int_0^t \mathbb{E}[\beta(s, X_s, \mathcal{L}(X_s | \mathcal{B}_s)) | \mathcal{B}_s] dB_s.$$

Let  $\overline{X}_t := X_t - \mathbb{E}[X_t | \mathcal{B}_t]$ , we have

$$\overline{X}_t = \overline{X}_0 + \int_0^t (b(s, X_s, \mathcal{L}(X_s | \mathcal{B}_s)) - \mathbb{E}[b(s, X_s, \mathcal{L}(X_s | \mathcal{B}_s)) | \mathcal{B}_s]) ds + \int_0^t \sigma(s, X_s, \mathcal{L}(X_s | \mathcal{B}_s)) dW_s.$$

Set  $N_t := \int_0^t \sigma(s, X_s, \mathcal{L}(X_s | \mathcal{B}_s)) dW_s$ , as an application of Itô's formula we have

$$\begin{aligned} \mathbb{E}[|N_t|^2 | \mathcal{B}_t] &= 2\mathbb{E}\left[\int_0^t N_s \cdot \sigma(s, X_s, \mathcal{L}(X_s | \mathcal{B}_s)) dW_s | \mathcal{B}_t\right] + \int_0^t \mathbb{E}[\text{Trace}(\sigma\sigma^\top)(s, X_s, \mathcal{L}(X_s | \mathcal{B}_s)) | \mathcal{B}_t] ds \\ &\leq C \int_0^t (1 + \overline{M}^2(\mathcal{L}(X_s | \mathcal{B}_s))) ds. \end{aligned}$$

By using Assumption 2 (i) also on the drift, we obtain

$$\overline{M}^2(\mathcal{L}(X_t | \mathcal{B}_t)) = \mathbb{E}[|\overline{X}|^2 | \mathcal{B}_t] \leq C \int_0^t (1 + \overline{M}^2(\mathcal{L}(X_s | \mathcal{B}_s))) ds.$$

Gronwall's lemma gives  $\overline{M}^2(\mathcal{L}(X_t | \mathcal{B}_t)) \leq C_T$ , where  $C_T > 0$  is a constant depending on  $T$ . A similar bound can be obtained for  $Z$ .

Combining the upper bounds on the central moment with Assumption 2, we have global Lipschitz continuity and boundedness of the coefficients  $b, \sigma, \beta$ . With standard estimates and Gronwall's lemma one can show that  $\mathbb{E}[\sup_{t \leq T} |X_t - Z_t|^2] = 0$ , which implies indistinguishability of the processes  $(X_t)_{t \in [0, T]}$  and  $(Z_t)_{t \in [0, T]}$ .  $\square$

*Remark 4.4.* Weak existence follows from [56] and one can apply Yamada-Watanabe to obtain well-posedness of (50). However, well-posedness will also follow from our results on mixed rough and stochastic McKean-Vlasov equations in the special case  $\beta(t, \pi) = P(\pi)$  defined in (60) below.

### 4.3 McKean-Vlasov with continuous deterministic forcing

Let  $b : \mathcal{P}_\rho(\mathbb{R}^D) \rightarrow \mathbb{R}^D$  and  $\sigma : \mathcal{P}_\rho(\mathbb{R}^D) \rightarrow \mathbb{R}^{D \times m}$  be measurable functions satisfying the following assumptions:

**Assumption 3.** Assume that there exists a constant  $C > 0$  such that

(i) (linear growth)  $\forall x \in \mathbb{R}^D, \forall \pi \in \mathcal{P}_\rho(\mathbb{R}^D)$ ,

$$|b(x, \pi)|, |\sigma(x, \pi)| \leq C(1 + \overline{M}^\rho(\pi))^{\frac{1}{\rho}}, \quad (51)$$

(ii) (Lipschitz continuity)  $\forall x, y \in \mathbb{R}^D, \forall \pi, \nu \in \mathcal{P}_\rho(\mathbb{R}^D)$ ,

$$|b(x, \pi) - b(y, \nu)|, |\sigma(x, \pi) - \sigma(y, \nu)| \leq C(1 + \overline{M}^\rho(\pi))^{\frac{1}{\rho}} (|x - y| + W_{\rho, \mathbb{R}^D}(\pi, \nu)). \quad (52)$$

Let  $F : [0, T] \rightarrow \mathbb{R}^D$  be a continuous bounded function,  $X_0 \in L_\omega^\rho$  and consider the following stochastic differential equation,

$$dX_t = b(X_t, \mathcal{L}(X_t)) dt + \sigma(X_t, \mathcal{L}(X_t)) dW_t + dF_t, \quad X_t|_{t=0} = X_0. \quad (53)$$

**Definition 4.5.** A stochastic process  $(X_t)_{t \in [0, T]}$  on  $\mathbb{R}^D$  is a solution for equation (53) with initial condition  $X_0$  if  $(\sigma(X_t, \mathcal{L}(X_t)))_{t \in [0, T]}$  is predictable and for every  $t \in [0, T]$ , the following integral equation is satisfied  $\mathbb{P}$ -a.s.,

$$X_t - F_t = \int_0^t b(X_s, \mathcal{L}(X_s)) ds + \int_0^t \sigma(X_s, \mathcal{L}(X_s)) dW_s, \quad X_t|_{t=0} = X_0.$$

*Remark 4.6.* In the following, we will construct an adapted solution  $(X_t)_{t \in [0, T]}$  with continuous sample paths. Since  $\sigma$  is Lipschitz continuous, we immediately have that  $(\sigma(X_t, \mathcal{L}(X_t)))_{t \in [0, T]}$  is predictable and the Itô integral is well defined.

Let  $X_t$  be a solution to equation (53), we define  $\overline{X}_t := X_t - \mathbb{E}[X_t]$ . We start with some preliminary expansions and estimates for  $\overline{X}$ . We have, for  $s, t \in [0, T]$ ,

$$\delta \overline{X}_{s,t} = \int_s^t (b(X_r, \mathcal{L}(X_r)) - \mathbb{E}[b(X_r, \mathcal{L}(X_r))]) dr + \int_s^t \sigma(X_r, \mathcal{L}(X_r)) dW_r. \quad (54)$$

**Lemma 4.7.** Let  $\rho \geq 2$  and  $X_0 \in L_\omega^\rho$ . There exists a constant  $C_T$  such that  $C_T \rightarrow 0$  as  $T \rightarrow 0$  and

$$\| \sup_{t \in [0, T]} |\overline{X}_t| \|_{L_\omega^\rho} \leq e^{C_T} [C_T + \|\overline{X}_0\|_{L_\omega^\rho}].$$

Moreover, for  $\alpha < \frac{1}{2}$  and  $\rho > \frac{2}{1-2\alpha}$ , we have

$$\|[\overline{X}]_\alpha\|_{L_\omega^\rho} \leq C_T(1 + \|\overline{X}_0\|_{L_\omega^\rho}).$$

*Proof.* Using standard estimates and the Burkholder-Davis-Gundy (BDG) inequality on equation (54) we have

$$\| \sup_{t \in [0, T]} |\overline{X}_t| \|_{L_\omega^\rho}^\rho \leq \|\overline{X}_0\|_{L_\omega^\rho}^\rho + C \int_0^T (1 + \overline{M}^\rho(\pi_t)) dt.$$

The first inequality follows from a standard application of Gronwall's lemma. For the second inequality we use again BDG and Jensen's inequality to obtain

$$\mathbb{E}[|\delta \overline{X}_{s,t}|^\rho] \leq |t - s|^{\rho/2 - \epsilon} C_T(1 + \|\overline{X}_0\|_{L_\omega^\rho}).$$

The second inequality as well as the condition on  $\rho$  follows from the Kolmogorov continuity theorem [48, Theorem 3.1].  $\square$

*Remark 4.8.* The bounds in Lemma 4.7 do not depend on the forcing term  $F$ . This gives us a good a-priori bound on the solution. Let  $\overline{T} > 0$  be fixed and arbitrarily large and assume that there exists  $M_0 > 0$  such that  $\|\overline{X}_0\|_{L^\rho} \leq M_0$ . Then there exists a global constant  $M(M_0, \overline{T})$  such that

$$\| \sup_{t \in [0, T]} |\overline{X}_t| \|_{L_\omega^\rho}, \|[\overline{X}]_\alpha\|_{L_\omega^\rho} \leq M. \quad (55)$$

We have the following a priori estimates.

**Lemma 4.9.** Let  $\rho \geq 2$ . Given  $X_0 \in L^\rho_\omega$  and  $F \in C_b([0, T], \mathbb{R}^D)$ , we call  $X(F, X_0)$  a solution to equation (53) with forcing  $F$  and initial condition  $X_0$ .

Let  $b, \sigma$  satisfy Assumption 3. Then there exists a constant  $C_T := C(T, \rho) > 0$  such that  $C_T \rightarrow 0$  as  $T \rightarrow 0$  and

$$\| \sup_{t \in [0, T]} |X_t(F, X_0)| \|_{L^\rho_\omega} \leq e^{C_T} [C_T + \|F\|_{C_b} + \|X_0\|_{L^\rho_\omega}]. \quad (56)$$

Given  $X_0, Y_0 \in L^\rho_\omega$  and  $F, G \in C_b([0, T], \mathbb{R}^D)$ , there exists a positive constant  $C > 0$  such that

$$\| \sup_{t \in [0, T]} |X_t(F, X_0) - X_t(G, Y_0)| \|_{L^\rho_\omega} \leq e^{C(1 + \|\bar{X}_0\|_{L^\rho_\omega})} \left( \|F - G\|_{C_b} + \|X_0 - Y_0\|_{L^\rho_\omega} \right). \quad (57)$$

*Proof.* We write  $X$  and omit here the dependence of the process on  $F$  and  $X_0$  as there is no possibility of confusion in the first part of the proof. Using Jensen's inequality, the Burkholder-Davis-Gundy inequality and Assumptions 3 we have the following estimate for  $s, t \in [0, T]$ ,

$$\begin{aligned} \mathbb{E} \sup_{t \in [0, T]} |X_t|^\rho &\leq \mathbb{E} |X_0|^\rho + CT^{\rho-1} \int_0^T \mathbb{E} |b(X_r, \mathcal{L}(X_r))|^\rho dr + CT^{\rho/2-1} \int_0^T \mathbb{E} |\sigma(X_r, \mathcal{L}(X_r))|^\rho dr + C \sup_{t \in [0, T]} |F_t|^\rho \\ &\leq \mathbb{E} |X_0|^\rho + C \left( T^{\rho-1} + T^{\rho/2-1} \right) \int_0^T (1 + \mathbb{E} |\bar{X}_r|^\rho) dr + C \sup_{t \in [0, T]} |F_t|^\rho, \end{aligned}$$

where the constant  $C$  depends only on  $\rho, b$  and  $\sigma$ . Equation (56) follows immediately using the bounds in Lemma 4.7.

We now prove inequality (57). Applying a similar reasoning as before, we obtain

$$\begin{aligned} \mathbb{E} \sup_{t \in [0, T]} |X_t(F, X_0) - X_t(G, Y_0)|^\rho &\leq C(T^\rho + T^{\rho/2}) \int_0^T (1 + \mathbb{E} |\bar{X}_t(F, X_0)|^\rho) \\ &\quad \cdot \left( \mathbb{E} \sup_{r \in [0, t]} |X_r(F, X_0) - X_r(G, Y_0)|^\rho + W_{\rho, \mathbb{R}^D}(\mathcal{L}(X_t(F, X_0)), \mathcal{L}(X_t(G, Y_0)))^\rho \right) dt + \|F - G\|_{C_b}^\rho. \end{aligned}$$

The  $\rho$ -Wasserstein distance is controlled by the  $L^\rho_\omega$  norm of the difference of the processes, so we can apply Gronwall's lemma and Lemma 4.7 to obtain the desired inequality (57).  $\square$

In addition to the previous bounds, if the forcing term is  $\alpha$ -Hölder continuous, we have that also the solution is  $\alpha$ -Hölder continuous.

**Lemma 4.10.** Let  $\alpha \in (\frac{1}{3}, \frac{1}{2})$  and  $\rho > 2/(1 - 2\alpha)$ . Assume that  $F, G \in C^\alpha([0, T], \mathbb{R}^D)$  and  $X_0, Y_0 \in L^\rho_\omega$ . Then there exist  $C_T, C := C(\rho, T, \alpha) > 0$  such that  $C_T \rightarrow 0$  as  $T \rightarrow 0$  and

$$\| [X(F, X_0)]_\alpha \|_{L^\rho_\omega} \leq C_T (1 + \|\bar{X}_0\|_{L^\rho_\omega}) + C[F]_\alpha. \quad (58)$$

Moreover,

$$\| [X(F, X_0) - X(G, Y_0)]_\alpha \|_{L^\rho_\omega} \leq C_T e^{C(1 + \|\bar{X}_0\|_{L^\rho_\omega})} \left( \|F - G\|_{C_b} + \|X_0 - Y_0\|_{L^\rho_\omega} \right) + C[F - G]_\alpha. \quad (59)$$

*Proof.* We write  $X$  and omit here the dependence of the process on  $F, X_0$  as there is no possibility of confusion in the first part of the proof. Using Jensen's inequality, the Burkholder-Davis-Gundy inequality and Assumption 3 we have the following estimate for  $s, t \in [0, T]$ ,

$$\begin{aligned} \mathbb{E} [|\delta(X - F)_{s,t}|^\rho] &\leq C|t - s|^{\rho-1} \int_s^t \mathbb{E} |b(X_r, \mathcal{L}(X_r))|^\rho dr + C|t - s|^{\rho/2-1} \int_s^t \mathbb{E} |\sigma(X_r, \mathcal{L}(X_r))|^\rho dr \\ &\leq C \left( |t - s|^{\rho-1} + |t - s|^{\rho/2-1} \right) \int_s^t (1 + \mathbb{E} |\bar{X}_r|^\rho) dr \\ &\leq C \left( |t - s|^{\rho-1} + |t - s|^{\rho/2-1} \right) |t - s| (1 + \mathbb{E} \sup_{r \in [0, T]} |\bar{X}_r|^\rho) \\ &\leq C|t - s|^{\rho/2-\epsilon} \left[ C_T (1 + \mathbb{E} \sup_{r \in [0, T]} |\bar{X}_r|^\rho) \right]. \end{aligned}$$

The constant  $C_T$  is such that  $C_T \rightarrow 0$ , as  $T \rightarrow 0$ . We apply Lemma 4.7 to obtain the following estimate,

$$\mathbb{E} [|\delta(X - F)_{s,t}|^\rho] \leq C_T |t - s|^{\rho/2-\epsilon} \left( 1 + \|\bar{X}_0\|_{L^\rho_\omega} \right).$$

Using the Kolmogorov continuity theorem we obtain that  $X - F$  is  $\alpha$ -Hölder continuous for  $\rho > 2/(1 - 2\alpha)$  and

$$\| [X]_\alpha \|_{L^\rho_\omega} \leq \| [X - F]_\alpha \|_{L^\rho_\omega} + C[F]_\alpha \leq C_T (1 + \|\bar{X}_0\|_{L^\rho_\omega}) + [F]_\alpha.$$

We now prove inequality (59). Arguing as in the first half of the proof we obtain

$$\begin{aligned} \mathbb{E}|\delta(X(F, X_0) - X(G, Y_0) - (F - G))_{s,t}|^\rho &\leq C \left( |t - s|^{\rho-1} + |t - s|^{\rho/2-1} \right) \\ &\cdot \int_s^t (1 + E|\bar{X}_r|^\rho) (\mathbb{E}|X_r(F, X_0) - X_r(G, Y_0)|^\rho + W_\rho(\mathcal{L}(X_r(F, X_0)), \mathcal{L}(X_r(G, Y_0)))^\rho) dr \end{aligned}$$

We apply Lemma 4.7 and (57) and use the fact that the  $L^\rho$ -norm of the difference of the process controls the  $\rho$ -Wasserstein distance, to obtain

$$\mathbb{E}|\delta(X(F, X_0) - X(G, Y_0) - F + G)_{s,t}|^\rho \leq C_T |t - s|^{\alpha\rho} e^{C_T(1 + \|\bar{X}_0\|_{L^\rho_\omega})} \left( \|F - G\|_{C_b} + \|X_0 - Y_0\|_{L^\rho_\omega} \right)^\rho.$$

Equation (59) follows as before from the Kolmogorov continuity theorem, for  $\rho > 2/(1 - 2\alpha)$ .  $\square$

**Lemma 4.11.** *Let  $F \in C_b([0, T], \mathbb{R}^D)$  and  $X_0 \in L^2_\omega$ . Let  $b$  and  $\sigma$  satisfy Assumption 3. Then equation (53) admits a unique strong solution.*

*Proof.* To prove existence we define

$$\tilde{b}(t, x, \mu) := b(x + F_t, (\tau_{F_t})_\# \mu), \quad t \in [0, T], x \in \mathbb{R}^D, \mu \in \mathcal{P}(\mathbb{R}^D),$$

where  $\tau_z : \mathbb{R}^D \rightarrow \mathbb{R}^D$  is the translation by  $z \in \mathbb{R}^D$ . Similarly we define  $\tilde{\sigma}$ . It is immediate to see that, if  $b, \sigma$  satisfy Assumption (3), then  $\tilde{b}, \tilde{\sigma}$  satisfy Assumption (1).

By Lemma 4.2 the following equation admits a unique global strong solution  $\tilde{X}$ ,

$$d\tilde{X}_t = \tilde{b}(t, \tilde{X}_t, \mathcal{L}(\tilde{X}_t)) dt + \tilde{\sigma}(t, \tilde{X}_t, \mathcal{L}(\tilde{X}_t)) dW_t, \quad \tilde{X}_t|_{t=0} = X_0 - F_0.$$

The stochastic process  $X = \tilde{X} + F$  solves equation (53). Uniqueness follows from the a priori estimates given in Lemma 4.9.  $\square$

## 4.4 Rough McKean-Vlasov

**Assumption 4.** *Let  $h \in C_b^2(\mathbb{R}^D, \mathbb{R}^d)$  such that  $x \mapsto xh^T(x) \in C_b^2(\mathbb{R}^D, \mathbb{R}^{D \times d})$ .*

For a given probability measure  $\pi \in \mathcal{P}(\mathbb{R}^D)$ , we define

$$P(\pi) := \text{Cov}_\pi(x, h)A + B, \quad \text{Cov}_\pi(x, h)^{l,j} = \int_{\mathbb{R}^D} (x^l - \pi[x^l])h^j(x)\pi(dx), \quad 1 \leq l \leq D, 1 \leq j \leq d, \quad (60)$$

where  $A \in \mathbb{R}^{d \times d}$  and  $B \in \mathbb{R}^{D \times d}$  are given matrices.

Let  $\mathbf{Y} := (Y, \mathbb{Y}) \in \mathcal{C}^\alpha(\mathbb{R}^d)$  and  $X_0 \in L^2_\omega$ . We study the following equation,

$$dX_t = b(X_t, \mathcal{L}(X_t)) dt + \sigma(X_t, \mathcal{L}(X_t)) dW_t + dF_t, \quad X_t|_{t=0} = X_0, \quad (61a)$$

$$dF_t = P(\mathcal{L}(X_t)) d\mathbf{Y}_t. \quad (61b)$$

**Definition 4.12.** A couple  $(X, F) : [0, T] \times \Omega \rightarrow \mathbb{R}^D \times \mathbb{R}^D$  is a solution to equation (61) if

- $F$  is a continuous path and  $X$  solves equation (61a) in the sense of Definition 4.5;
- $P(\mathcal{L}(X_t)) \in \mathcal{D}_{\mathbf{Y}}^{2\alpha}([0, T], \mathcal{L}(\mathbb{R}^d, \mathbb{R}^D))$  and  $F$  is the rough integral in (61b).

Next we prove that, if  $X$  solves (61a) for a fixed controlled  $F$ , then  $P(\mathcal{L}(X_t))$  is a controlled path, which makes the rough integral (61b) well defined.

**Lemma 4.13.** *Let  $\alpha \in (\frac{1}{3}, \frac{1}{2})$  and  $\rho > 2/(1 - 2\alpha)$ . Let  $\mathbf{Y} := (Y, \mathbb{Y}) \in \mathcal{C}^\alpha(\mathbb{R}^d)$ ,  $F \in \mathcal{D}_{\mathbf{Y}}^{2\alpha}([0, T], \mathbb{R}^D)$  and  $X_0 \in L^\rho_\omega(\mathbb{R}^D)$ .*

*If  $\pi_t$  is the law of the solution process  $X_t(F, X_0)$  to equation (53) with forcing  $F$  and initial condition  $X_0$ , then*

$$P(\pi_\cdot) \in \mathcal{D}_{\mathbf{Y}}^{2\alpha}([0, T], \mathcal{L}(\mathbb{R}^d, \mathbb{R}^D)),$$

*with Gubinelli derivative*

$$P(\pi_s)' = (F_s')^\top \pi[(x - \pi[x])Dh^\top]A. \quad (62)$$

*Moreover, we have the bound*

$$\|P(\pi_\cdot)\|_{\mathcal{D}_{\mathbf{Y}}^{2\alpha}} \leq C \|h\|_{C_b^2} (1 + \|\bar{X}_0\|_{L^\rho_\omega} + [F]_\alpha) \left( C_T(1 + \|\bar{X}_0\|_{L^\rho_\omega}) + \|F'\|_{C_b}[Y]_\alpha + [R^F]_{2\alpha} \right),$$

*where  $C_T \rightarrow 0$  as  $T \rightarrow 0$ .*

*Proof.* To simplify the notation, we write  $X_t$  for  $X(F, X_0)$ . Let  $\bar{X}_t := X_t - \mathbb{E}[X_t]$ . We have the following expansion for  $h(X)$ ,  $s, t \in [0, T]$ ,

$$\begin{aligned} \delta h(X)_{s,t} = & [Dh]_{s,t}^{1,x} \left( \int_s^t b(X_r, \mathcal{L}(X_r)) dr + R_{s,t}^F \right) \\ & + ([D^2 h]_{s,t}^{2,x} \delta X_{s,t} + Dh(X_s)) \left( \int_s^t \sigma(X_r, \mathcal{L}(X_r)) dW_r + F'_s \delta Y_{s,t} \right). \end{aligned} \quad (63)$$

Let  $s, t \in [0, T]$ . From the definition of  $P(\pi)$ , equation (60), we have

$$\delta P(\pi)_{s,t} = \delta \mathbb{E}[\bar{X} \cdot h(X)]_{s,t}^\top A = \mathbb{E}[\delta \bar{X}_{s,t} h(X_s)^\top] A + \mathbb{E}[\delta \bar{X}_{s,t} \delta h(X)_{s,t}^\top] A + \mathbb{E}[\bar{X}_s \delta h(X)_{s,t}^\top] A =: I_1 + I_2 + I_3.$$

We expand  $I_3$  even further using (63)

$$\begin{aligned} I_3 = & \mathbb{E}[\bar{X}_s (Dh(X_s) F'_s \delta Y_{s,t})^\top] A + \mathbb{E} \left[ \bar{X}_s \left( Dh(X_s) \int_s^t \sigma(X_r, \mathcal{L}(X_r)) dW_r \right)^\top \right] A \\ & + \mathbb{E} \left[ \bar{X}_s \left( [Dh]_{s,t}^{1,x} \left( \int_s^t b(X_r, \mathcal{L}(X_r)) dr + R_{s,t}^F \right) + [D^2 h]_{s,t}^{2,x} \delta X_{s,t} \left( \int_s^t \sigma(X_r, \mathcal{L}(X_r)) dW_r + F'_s \delta Y_{s,t} \right) \right)^\top \right] A \\ = & : \mathbb{E} \left[ \bar{X}_s (Dh(X_s) F'_s \delta Y_{s,t})^\top \right] A + I_4. \end{aligned}$$

We can write

$$\delta P(\pi)_{s,t} = \mathbb{E}[\bar{X}_s (Dh(X_s) F'_s \delta Y_{s,t})^\top] A + R_{s,t}^P,$$

where  $R_{s,t}^P = I_1 + I_2 + I_4$ . For  $1 \leq l \leq D$  and  $1 \leq j \leq d$  we write the  $(l, j)$  entry of the matrix  $P$  as

$$\delta P^{l,j}(\pi)_{s,t} = (P^{l,j}(\pi))'_s \cdot \delta Y_{s,t} + R_{s,t}^{P^{l,j}},$$

where, for  $s \in [0, T]$ ,  $P(\pi_s)' \in \mathcal{L}(\mathbb{R}^d, \mathcal{L}(\mathbb{R}^d, \mathbb{R}^D)) \cong \mathbb{R}^{D \times d \times d}$  is the Gubinelli derivative given as

$$(P^{l,j}(\pi_s))' := \sum_{k=1}^D \mathbb{E}[\bar{X}_s^l (Dh(X_s)^\top A)^j] (F'_s)^k = \sum_{k=1}^D \sum_{i=1}^d \mathbb{E}[\bar{X}_s^l \partial_k h^i(X_s)] A^{i,j} (F'_s)^k \in \mathbb{R}^d.$$

We check now that  $R^P$  is a remainder with regularity  $2\alpha$ . Using the martingale property of the stochastic integral in (54), we have

$$I_1 = \mathbb{E}[\mathbb{E}_s[\delta \bar{X}_{s,t}] h(X_s)^\top] A = \mathbb{E} \left[ \int_s^t (b(X_r, \mathcal{L}(X_r)) - \mathbb{E}[b(X_r, \mathcal{L}(X_r))]) dr h(X_s)^\top \right] A.$$

We apply Lemma 4.7 to obtain

$$|I_1| \leq |t - s| C \|h\|_\infty (C_T + \|X_0\|_{L_\omega^\rho}),$$

where  $C > 0$  is a global constant. For  $I_2$  we use Lemma 4.7 and (58) to obtain, for any  $\rho > 3$ ,

$$\begin{aligned} |I_2| = & |\mathbb{E}[\delta \bar{X}_{s,t} ([Dh]_{s,t}^{1,x} \delta X_{s,t})^\top]| \leq \|Dh\|_\infty \|\bar{X}\|_\alpha \| [X]_\alpha \|_{L_\omega^\rho} \\ & \leq \|Dh\|_\infty C_T (1 + \|\bar{X}_0\|_{L_\omega^\rho}) [C_T (1 + \|\bar{X}_0\|_{L_\omega^\rho}) + [F]_\alpha] |t - s|^{2\alpha}. \end{aligned}$$

We proceed by finding a bound for  $I_4$ . Notice that the term  $\mathbb{E}[\bar{X}_s (Dh(X_s) \int_s^t \sigma(X_r, \mathcal{L}(X_r)) dW_r)^\top]$  vanishes thanks to the martingale property of the stochastic integral. Using Lemma 4.7 and inequality (58), we get

$$\begin{aligned} I_4 \leq & C |t - s|^{2\alpha} \|Dh\|_\infty \sup_{t \in [0, T]} \|\bar{X}_t\|_{L_\omega^\rho} \left( C_T (1 + \sup_{t \in [0, T]} \|\bar{X}_t\|_{L_\omega^\rho}) + [R^F]_{2\alpha} \right) \\ & + C |t - s|^{2\alpha} \|D^2 h\|_\infty \|[X]_\alpha\|_{L_\omega^\rho} \left( C_T (1 + \sup_{t \in [0, T]} \|\bar{X}_t\|_{L_\omega^\rho}) + \|F'\|_{C_b[Y]_\alpha} \right) \\ \leq & C |t - s|^{2\alpha} \|Dh\|_\infty (C_T + \|\bar{X}_0\|_{L_\omega^\rho}) \left( C_T (1 + \|\bar{X}_0\|_{L_\omega^\rho}) + [R^F]_{2\alpha} \right) \\ & + C |t - s|^{2\alpha} \|D^2 h\|_\infty (1 + \|\bar{X}_0\|_{L_\omega^\rho} + [F]_\alpha) \left( C_T (1 + \|\bar{X}_0\|_{L_\omega^\rho}) + \|F'\|_{C_b[Y]_\alpha} \right) \\ \leq & C |t - s|^{2\alpha} (\|Dh\|_\infty + \|D^2 h\|_\infty) (1 + \|\bar{X}_0\|_{L_\omega^\rho} + [F]_\alpha) \left( C_T (1 + \|\bar{X}_0\|_{L_\omega^\rho}) + \|F'\|_{C_b[Y]_\alpha} + [R^F]_{2\alpha} \right). \end{aligned}$$

□

We now set up a contraction argument that we will use to prove the well-posedness of equation (61). Let  $f \in \mathcal{D}_{\mathbf{Y}}^{2\alpha}([0, T], \mathcal{L}(\mathbb{R}^d, \mathbb{R}^D))$ . We define, for  $t \in [0, T]$ ,

$$(F_t, F'_t) := \left( \int_0^t f_r d\mathbf{Y}_r, f_t \right) \in \mathcal{D}_{\mathbf{Y}}^{2\alpha}([0, T], \mathbb{R}^D). \quad (64)$$

Now we plug this as the forcing into equation (53) with initial condition  $X_0 \in L_\omega^\rho$  and call  $X$  the solution with law  $\pi$ . We define the map

$$\begin{aligned} \Gamma : \mathcal{D}_{\mathbf{Y}}^{2\alpha}([0, T], \mathcal{L}(\mathbb{R}^d, \mathbb{R}^D)) &\rightarrow \mathcal{D}_{\mathbf{Y}}^{2\alpha}([0, T], \mathcal{L}(\mathbb{R}^d, \mathbb{R}^D)) \\ f &\mapsto P(\pi.). \end{aligned}$$

From [48] we have the following estimates,

$$\|f\|_{C_b}, \|f'\|_{C^b} \leq |f_0| + \|f\|_{\mathcal{D}_{\mathbf{Y}}^{2\alpha}},$$

$$[F]_\alpha, [R^F]_\alpha \leq C \|\mathbf{Y}\|_\alpha (|f_0| + \|f\|_{\mathcal{D}_{\mathbf{Y}}^{2\alpha}}).$$

It follows from Lemma 4.13 that  $\Gamma$  is well defined. Let us prove that there exists a small time, such that the following set is invariant under  $\Gamma$ ,

$$D(\Gamma) := \{f \in \mathcal{D}_{\mathbf{Y}}^{2\alpha} \mid |f_0| + \|f\|_{\mathcal{D}_{\mathbf{Y}}^{2\alpha}} \leq 2\|h\|_\infty M\},$$

where  $M$  is the global constant of Remark 4.8.

We have the following estimates,

$$|\Gamma(f)_0| = |\mathbb{E}[\bar{X}_0 h(X_0)]A + B| \leq \|h\|_\infty M,$$

$$\begin{aligned} \|\Gamma(f)\|_{\mathcal{D}_{\mathbf{Y}}^{2\alpha}} &= \|P(\pi.)\|_{\mathcal{D}_{\mathbf{Y}}^{2\alpha}} \leq C \|h\|_{C_b^2} (1 + \|\bar{X}_0\|_{L_\omega^\rho} + \|\mathbf{Y}\|_\alpha (|f_0| + \|f\|_{\mathcal{D}_{\mathbf{Y}}^{2\alpha}})) \left( C_T (1 + \|\bar{X}_0\|_{L_\omega^\rho}) + \|\mathbf{Y}\|_\alpha (|f_0| + \|f\|_{\mathcal{D}_{\mathbf{Y}}^{2\alpha}}) \right) \\ &\leq \|h\|_{C_b^2} C_T (1 + M + 2\|h\|_\infty (\|\mathbf{Y}\|_\alpha \vee \|\mathbf{Y}\|_{\bar{\alpha}}))^2 \\ &\leq \|h\|_\infty M, \end{aligned}$$

where in the last step we chose  $T$  small enough such that

$$C_T \leq \frac{\|h\|_\infty M}{\|h\|_{C_b^2} (1 + M + 2M\|h\|_\infty (\|\mathbf{Y}\|_\alpha \vee \|\mathbf{Y}\|_{\bar{\alpha}}))^2}.$$

Adding the previous estimates we proved that  $\Gamma(D(\Gamma)) \subset D(\Gamma)$ . Since  $C_T$  only depends on global quantities, we can divide each interval  $[0, \bar{T}]$  into smaller intervals of length  $T$  and in each one apply the contraction argument. We prove in the following that  $\Gamma$  is a contraction on  $B_T$ . We start by showing that  $\Gamma$  is Lipschitz continuous.

**Lemma 4.14.** *Let  $1/3 < \alpha < \bar{\alpha} < 1/2$ . Let  $\rho > 2/(1 - 2\alpha)$  and  $X_0, Z_0 \in L_\omega^\rho$ . Let  $\mathbf{Y}, \hat{\mathbf{Y}} \in \mathcal{C}^{\bar{\alpha}}$  and  $F \in \mathcal{D}_{\mathbf{Y}}^{2\alpha}$ ,  $G \in \mathcal{D}_{\hat{\mathbf{Y}}}^{2\alpha}$ . Assume that there exists a universal constant  $C > 0$  such that*

$$\|F\|_{\mathcal{D}_{\mathbf{Y}}^{2\alpha}} \leq C \|\mathbf{Y}\|_\alpha, \quad \|G\|_{\mathcal{D}_{\hat{\mathbf{Y}}}^{2\alpha}} \leq C \|\bar{\mathbf{Y}}\|_\alpha, \quad \|\bar{X}_0\|_{L_\omega^\rho}, \|\bar{Z}_0\|_{L_\omega^\rho} \leq C.$$

*We call  $X_t(F, X_0)$  (resp.  $X_t(G, Z_0)$ ) the solution to equation (53) with inputs  $F$  and  $X_0$  (resp.  $G$  and  $Z_0$ ). We call  $R$  the difference of the remainders of  $P(\pi^x)$  and  $P(\pi^z)$ . There exists a global constant  $C_T > 0$  such that  $C_T \rightarrow 0$  as  $T \rightarrow 0$  and*

$$[R]_{2\alpha} \leq C_T \left( \|F - G\|_{C_b} + \|X_0 - Z_0\|_{L_\omega^\rho} + [F - G]_\alpha \right) + \|F' - G'\|_{C_b} [\mathbf{Y}]_\alpha + \|G'\|_{C_b} [\mathbf{Y} - \hat{\mathbf{Y}}]_\alpha. \quad (65)$$

*A similar estimate can be obtained for the difference of the Gubinelli derivatives.*

*Proof.* To simplify the notation, we write  $X_t$  for  $X(F, X_0)_t$  and  $Z_t$  for  $X(G, Z_0)_t$ . Let  $\bar{X}_t := X_t - E[X_t]$ ,

similarly we define  $\bar{Z}_t$ . We have the following expansion for  $h(X) - h(Z)$ ,  $s, t \in [0, T]$ ,  $\mathbb{P}$ -a.s.,

$$\delta(h(X_\cdot) - h(Z_\cdot))_{s,t} = Dh(X_s)F'\delta Y_{s,t} - Dh(X_s)G'\delta \hat{Y}_{s,t} \quad (66a)$$

$$+ Dh(X_s) \left( \int_s^t \sigma(X_r, \mathcal{L}(X_r)) dW_r \right) - Dh(Z_s) \left( \int_s^t \sigma(Z_r, \mathcal{L}(Z_r)) dW_r \right) \quad (66b)$$

$$+ [Dh]_{s,t}^{1,z} \left( \int_s^t [b(X_r, \mathcal{L}(X_r)) - b(Z_r, \mathcal{L}(Z_r))] dr + R_{s,t}^F - R_{s,t}^G \right) \quad (66c)$$

$$+ ([Dh]_{s,t}^{1,x} - [Dh]_{s,t}^{1,z}) \left( \int_s^t b(X_r, \mathcal{L}(X_r)) dr + R_{s,t}^F \right) \quad (66d)$$

$$+ [D^2 h]_{s,t}^{2,z} \delta Z_{s,t} \left( \int_s^t [\sigma(X_r, \mathcal{L}(X_r)) - \sigma(Z_r, \mathcal{L}(Z_r))] dW_r \right) \quad (66e)$$

$$+ [D^2 h]_{s,t}^{2,z} \delta Z_{s,t} (F'_s \delta Y_{s,t} - G'_s \delta \hat{Y}_{s,t}) \quad (66f)$$

$$+ ([D^2 h]_{s,t}^{2,z} \delta(X - Z)_{s,t} + ([D^2 h]_{s,t}^{2,x} - [D^2 h]_{s,t}^{2,z}) \delta X_{s,t}) \left( \int_s^t \sigma(X_r, \mathcal{L}(X_r)) dW_r \right). \quad (66g)$$

Let  $s, t \in [0, T]$ . We call  $\pi^x := \mathcal{L}(X)$  and  $\pi^z := \mathcal{L}(Z)$ . From the definition of  $P(\pi)$ , equation (60), we have

$$\begin{aligned} \delta(P(\pi^x) - P(\pi^z))_{s,t} &= \mathbb{E}[\delta(\bar{X}_\cdot - \bar{Z}_\cdot)_{s,t} h(Z_s)^T] A + \mathbb{E}[\delta(\bar{X}_\cdot - \bar{Z}_\cdot)_{s,t} \delta h(Z_\cdot)_{s,t}^T] A + \mathbb{E}[(\bar{X}_s - \bar{Z}_s) \delta h(Z_\cdot)_{s,t}^T] A \\ &\quad + \mathbb{E}[\delta \bar{X}_{s,t} (h(X_s) - h(Z_s))] A + \mathbb{E}[\delta \bar{X}_{s,t} \delta(h(X_\cdot) - h(Z_\cdot))_{s,t}] A + \mathbb{E}[\bar{X}_s \delta(h(X_\cdot) - h(Z_\cdot))_{s,t}] A \\ &=: I_{11} + I_{12} + I_{13} + I_{21} + I_{22} + I_{23}. \end{aligned}$$

Using the same expansion for  $h(Z)$  as in (63), we obtain

$$\begin{aligned} I_{13} &= \mathbb{E}[(\bar{X}_s - \bar{Z}_s) (Dh(Z_s) G' \delta \hat{Y}_{s,t})^T] A + 0 + \mathbb{E}[(\bar{X}_s - \bar{Z}_s) ([Dh]_{s,t}^{1,z} (\int_s^t b(Z_r, \mathcal{L}(Z_r)) dr + R_{s,t}^G))^T] A \\ &\quad + \mathbb{E}[(\bar{X}_s - \bar{Z}_s) ([D^2]_{s,t}^{2,z} \delta Z_{s,t} (\int_s^t \sigma(Z_r, \mathcal{L}(Z_r)) dW_r + G'_s \delta \hat{Y}_{s,t}))^T] A \\ &= \mathbb{E}[(\bar{X}_s - \bar{Z}_s) (Dh(Z_s) G' \delta \hat{Y}_{s,t})^T] A + I_{14} + I_{15}. \end{aligned}$$

The second term in the first line vanishes thanks to the martingale property of the stochastic integral. Similarly, we expand  $I_{23}$  using (66). Notice that the term (66b) vanishes because the stochastic integral is a martingale. The term (66a) produces a term of regularity  $\alpha$ , the others have regularity  $2\alpha$ .

$$I_{23} = \mathbb{E}[\bar{X}_s (Dh(X_s) F' \delta Y_{s,t} - Dh(Z_s) G' \delta \hat{Y}_{s,t})^T] A + I_{24},$$

where  $I_{24} := \mathbb{E}[\bar{X}_s ((66c) + \dots + (66g))] A$ . We thus obtain that

$$\delta(P(\pi^x) - P(\pi^z))_{s,t} = \mathbb{E}[(\bar{X}_s - \bar{Z}_s) (Dh(Z_s) G' \delta \hat{Y}_{s,t})^T] A + \mathbb{E}[\bar{X}_s (Dh(X_s) F' \delta Y_{s,t} - Dh(X_s) G' \delta \hat{Y}_{s,t})^T] A + R_{s,t}$$

with  $R_{s,t} = I_{11} + I_{12} + I_{14} + I_{15} + I_{21} + I_{22} + I_{24}$ . For  $1 \leq l \leq D$  and  $1 \leq j \leq d$  we write the  $(l, j)$  entry of the matrix as

$$\begin{aligned} \delta(P(\pi^x) - P(\pi^z))_{s,t}^{l,j} &= \sum_{k=1}^D \left( \mathbb{E}[(\bar{X}_s - \bar{Z}_s)^l (Dh(Z_s)^\top A)^{i,k} (G')^k] + \mathbb{E}[(\bar{X}_s - \bar{Z}_s) ((Dh(X_s) - Dh(Z_s))^\top A)^{i,k} (G')^k] \right. \\ &\quad \left. + \mathbb{E}[\bar{X}_s (Dh(X_s)^\top A)^{i,k} (F' - G')^k] \right) \cdot \delta \hat{Y}_{s,t} + \sum_{k=1}^D \mathbb{E}[\bar{X}_s^l (Dh(X_s)^\top A)^{i,k} (F')^k] \cdot \delta(Y - \hat{Y})_{s,t} + R_{s,t}^{P^{l,j}}. \end{aligned}$$

We must now find estimates for  $[R]_{2\alpha}$ . We start by some estimates of the processes in  $L^\rho$ . From equation (57) we have

$$\| \sup_{t \in [0, T]} |X_t - Z_t| \|_{L_\omega^\rho} \leq C \left( \|F - G\|_{C_b} + \|X_0 - Y_0\|_{L_\omega^\rho} \right). \quad (67)$$

It follows from the equations for  $\bar{X}$  and  $\bar{Z}$  as well as estimate (67) that

$$\|[\bar{X} - \bar{Z}]_\alpha\|_{L_\omega^\rho} \leq C_T (1 + \|\bar{X}_0\|_{L_\omega^\rho}) \sup_{t \in [0, T]} \|X_t - Z_t\|_{L_\omega^\rho} \leq C_T \left( \|F - G\|_{C_b} + \|X_0 - Y_0\|_{L_\omega^\rho} \right).$$

Using Lemma 4.7 we also have

$$\|[\bar{X} - \bar{Z}]_\alpha\|_{L_\omega^\rho} \leq \|[\bar{X}]_\alpha\|_{L_\omega^\rho} + \|[\bar{Z}]_\alpha\|_{L_\omega^\rho} \leq C_T.$$

From (59) we have

$$\| [X - Z]_\alpha \|_{L^\rho_\omega} \leq C_T \left( \|F - G\|_{C_b} + \|X_0 - Y_0\|_{L^\rho_\omega} \right) + C[F - G]_\alpha.$$

From (58) we obtain

$$\| [X]_\alpha \|_{L^\rho_\omega}, \| [Z]_\alpha \|_{L^\rho_\omega} \leq C_T.$$

Using the previous estimates we obtain

$$\begin{aligned} \delta(I_{11} + I_{21})_{s,t} &\leq C_T \|h\|_{C_b^1} \sup_{t \in [0,T]} |X_t - Z_t| \|L^\rho_\omega |t - s|^{2\alpha}, \\ \delta(I_{12} + I_{22})_{s,t} &\leq C_T \|h\|_{C_b^1} \| [X - Z]_\alpha \|_{L^\rho_\omega} |t - s|^{2\alpha}, \\ \delta(I_{14} + I_{15})_{s,t} &\leq \|h\|_{C_b^2} \left[ \sup_{t \in [0,T]} |X_t - Z_t| \|L^\rho_\omega (C_T + [R^G]_{2\alpha} + \|G'\|_{C_b} [\hat{Y}]_\alpha) |t - s|^{2\alpha} \right]. \end{aligned}$$

For  $I_{24}$  we have a combination of the above and an extra term

$$[I_{24}]_{2\alpha} \leq \dots + \|F' - G'\|_{C_b} [\mathbf{Y}]_\alpha + \|G'\|_{C_b} [\mathbf{Y} - \hat{\mathbf{Y}}]_\alpha.$$

Summing up, we get

$$[R]_{2\alpha} \leq C \left( \|F - G\|_{C_b} + \|X_0 - Y_0\|_{L^\rho_\omega} + [F - G]_\alpha \right) (C_T + [R^G]_{2\alpha} + \|G'\|_{C_b} [\hat{Y}]_\alpha) + \|F' - G'\|_{C_b} [\mathbf{Y}]_\alpha + \|G'\|_{C_b} [\mathbf{Y} - \hat{\mathbf{Y}}]_\alpha,$$

which concludes the proof using the assumption  $\|G\|_{\mathcal{D}^{2\alpha}_{\mathbf{Y}}} \leq C \|\bar{\mathbf{Y}}\|_\alpha$   $\square$

We are now ready to prove the main well-posedness result.

**Theorem 4.15.** *Let  $1/3 < \alpha < \bar{\alpha} < 1/2$ . Let  $\rho > 2/(1 - 2\alpha)$  and  $X_0, Z_0 \in L^\rho$ . Let  $X_0 \in L^\rho$  and  $\mathbf{Y} \in \mathcal{C}^{\bar{\alpha}}(\mathbb{R}^d)$ . If  $b, \sigma$  and  $h$  satisfy Assumptions 3 and 4, then equation (61) admits a unique solution  $(X, F) \in L^\rho(C([0, T], \mathbb{R}^D)) \times \mathcal{D}^{2\alpha}_{\mathbf{Y}}([0, T], \mathbb{R}^D)$ .*

*Proof.* A process  $X : \Omega \rightarrow C([0, T], \mathbb{R}^D)$  is a solution to equation (61) if and only if  $P(\mathcal{L}(X))$  is a fixed point of  $\Gamma$ .

We want to prove that  $\Gamma : D(\Gamma) \rightarrow D(\Gamma)$  is a contraction. If  $f, g \in D(\Gamma)$  we have that  $F, G$  defined as in (64) satisfy the assumptions of Lemma 4.14 with a global constant  $C = C(h, M)$ . By taking  $\mathbf{Y} = \hat{\mathbf{Y}}$  and  $X_0 = Z_0$  the estimate in Lemma 4.14 reduces to

$$\|\Gamma(f) - \Gamma(g)\|_{\mathcal{D}^{2\alpha}_{\mathbf{Y}}} \leq C_T(M, \|\mathbf{Y}\|_{\bar{\alpha}}) (\|f - g\|_{\mathcal{D}^{2\alpha}_{\mathbf{Y}}} - \|f_0 - g_0\|_{C_b}), \quad f, g \in D(\Gamma).$$

Choosing now  $T_0$  small enough such that  $C_{T_0} < 1$ , we have that  $\Gamma$  is a contraction on the closed subset  $D(\Gamma) \subset \mathcal{D}^{2\alpha}_{\mathbf{Y}}([0, T_0], \mathcal{L}(\mathbb{R}^d, \mathbb{R}^D))$ . Hence,  $\Gamma$  admits a unique fixed point.

Since the small constant  $C_{T_0}$  in the definition of the domain  $D(\Gamma)$  of  $\Gamma$  and in the contraction argument only depends on the global quantities  $h, M, \mathbf{Y}$  and not on the initial condition  $X_0$ , we can construct a finite family of subsequent time intervals of size  $T_0$  that covers  $[0, T]$ . On each of these time interval we construct the solution as a fixed point of  $\Gamma$ .  $\square$

**Corollary 4.16.** *Let  $1/3 < \bar{\alpha} < 1/2$  and  $\rho > 2/(1 - 2\bar{\alpha})$ . Let  $X_0 \in L^\rho$  and  $\mathbf{Y}^1, \mathbf{Y}^2 \in \mathcal{C}^{\bar{\alpha}}$ . For  $i = 1, 2$ , let  $X^i$  be the solution to equation (61) with driver  $\mathbf{Y}^i$  and initial condition  $X_0$ . Call  $\pi^i := \mathcal{L}(X^i) \in \mathcal{P}_\rho(C([0, T], \mathbb{R}^D))$ . There exists a positive constant  $C > 0$  such that*

$$W_{\rho, C([0, T], \mathbb{R}^D)}(\pi^1, \pi^2) \leq C \rho_{\bar{\alpha}}(\mathbf{Y}^1, \mathbf{Y}^2).$$

*Proof.* It follows from Lemma 4.9 and Lemma 4.14 that the solution map  $\mathbf{Y} \mapsto X$  of equation (61) is a Lipschitz-continuous function between the spaces  $\mathcal{C}^{\bar{\alpha}}(\mathbb{R}^d)$  and  $L^\rho(C([0, T], \mathbb{R}^D))$ . The corollary follows as the Wasserstein distance between  $\pi^1$  and  $\pi^2$  is always less or equal to the  $L^\rho$  distance between  $X^1$  and  $X^2$ .  $\square$

We now proceed to prove a Wong-Zakai type result when  $\mathbf{Y}$  is the Itô lift of a Brownian motion. We introduce the approximation  $X^n$  as the solution of

$$dX_t^n = b(X_t^n, \mathcal{L}(X_t^n)) dt + \sigma(X_t^n, \mathcal{L}(X_t^n)) dW_t + dF_t^n, \quad X_t|_{t=0} = X_0, \quad (68a)$$

$$\dot{F}_t^n = P(\mathcal{L}(X_t^n)) \dot{Y}_t^n + \frac{1}{2} P(\mathcal{L}(X_t^n))^\top \mathbb{E}[\bar{X}_t^n D h(X_t^n)^\top] A. \quad (68b)$$

where  $Y^n$  is a piecewise linear approximation of  $Y$ . We have the following result.

**Theorem 4.17.** *The solution  $X^n$  converges to  $X$  in the Wasserstein distance, viz  $\mathbb{P}$ -a.s. we have*

$$W_{\rho, C([0, T], \mathbb{R}^D)}(\mathcal{L}(X^n), \mathcal{L}(X)) \rightarrow 0$$

as  $n \rightarrow \infty$ .

*Proof.* We start by noting that  $Y^n$  solves (68) if and only if it solves the rough path equation

$$\begin{aligned} dX_t^n &= b(X_t^n, \mathcal{L}(X_t^n)) dt + \sigma(X_t^n, \mathcal{L}(X_t^n)) dW_t + dF_t^n, & X_t|_{t=0} &= X_0, \\ dF_t^n &= P(\mathcal{L}(X_t^n)) d\mathbf{Y}_t^{n,Ito}, \end{aligned}$$

where

$$\mathbf{Y}_{st}^{n,Ito} = \left( \delta Y_{s,t}^n, \int_s^t \delta Y_{s,r}^n \otimes \dot{Y}_r^n dr + \frac{1}{2}(t-s)I_{d \times d} \right).$$

It is well known that the canonical lift of  $Y^n$ ,

$$\mathbf{Y}_{st}^{n,Str} = \left( \delta Y_{s,t}^n, \int_s^t \delta Y_{s,r}^n \otimes \dot{Y}_r^n dr \right),$$

converges  $\mathbb{P}$ -a.s. in the rough path topology to the rough path

$$\mathbf{Y}_{st}^{Str} = \left( \delta Y_{s,t}, \int_s^t \delta Y_{s,r} \otimes \odot dY_r \right).$$

where the latter integral is the Stratonovich integral. From this it follows immediately that

$$\mathbf{Y}_{st}^{n,Ito} \rightarrow \left( \delta Y_{s,t}, \int_s^t \delta Y_{s,r} \otimes \odot dY_r + \frac{1}{2}(t-s)I_{d \times d} \right) = \left( \delta Y_{s,t}, \int_s^t \delta Y_{s,r} \otimes dY_r \right)$$

where the latter integral is the Itô integral. The result now follows from Corollary 4.16.  $\square$

## 4.5 Proof of Theorem 1.1

We can now proceed with the proof of the main theorem.

**Well-posedness of the rough stochastic McKean-Vlasov equation (11).** Given the coefficients of equation (11) we transform it into equation (61) by defining the following coefficients

$$\begin{aligned} b(x, \pi) &:= f(x) - P(\pi)h(x), \\ \sigma(x, \pi) &:= \left( G^{\frac{1}{2}} - P(\pi)U, \quad -P(\pi)R^{\frac{1}{2}} \right), \\ P(\pi) &:= \text{Cov}_\pi(x, h)C^{-1} + B. \end{aligned}$$

The  $m$ -Brownian motion  $W$  in (61) stands for the paired independent Brownian Motions  $(\widehat{W}, \widehat{V})$  of equation (11), with  $m = D + d$ . Moreover, if  $\mathbf{Y} = (Y, \mathbb{Y})$  is any given rough path, we can modify it by a bounded variation term to obtain

$$\overline{\mathbf{Y}}_{s,t} := (Y_{s,t}, \mathbb{Y}_{s,t} + \frac{1}{2}(t-s)I_{d \times d}).$$

We have that equation (61) driven by  $\overline{\mathbf{Y}}$  corresponds to equation (11) driven by  $\mathbf{Y}$  and the term  $\frac{1}{2}(t-s)I_{d \times d}$  generates the correction term  $\Gamma(\pi)$ . Hence, the proof of well-posedness and stability of equation (11) follows from Theorem 4.15 and Corollary 4.16.

**Well-posedness of the McKean-Vlasov equation with common noise (10).** The uniqueness of equation (10) follows from Lemma 4.3, as the coefficients of equation (10) satisfy Assumption 2. The following lemma gives existence.

**Lemma 4.18.** *Let  $X^{\mathbf{Y}}$  denote the solution of (11) and let  $\mathbf{Y}(\omega)$  denote the Stratonovich rough path lift of the Brownian motion  $Y$ . Then  $X^{\mathbf{Y}}$  is the solution of (10) and we have*

$$\mathbb{E} \left[ \phi(X_t^{\mathbf{Y}}) | \mathcal{Y}_t \right] (\omega) = \mathbb{E} \left[ \phi(X_t^{\mathbf{Y}}) \right] |_{\mathbf{Y}=\mathbf{Y}(\omega)|_{[0,t]}}, \quad \mathbb{P}_Y - a.e. \omega. \quad (69)$$

*Proof.* Since the Stratonovich rough path lift of  $Y|_{[0,t]}$  coincides with  $(Y, \mathbb{Y})|_{[0,t]}$ , we notice that by uniqueness, we have

$$X_t^{\mathbf{Y}} = X_t^{\mathbf{Y}|_{[0,t]}}.$$

Since  $\mathbf{Y}|_{[0,t]}$  is independent of  $\sigma(W_s, X_0, 0 \leq s \leq t)$ , then it is also independent of  $X_t^{\mathbf{Y}}$  for every  $\mathbf{y} \in \mathcal{C}^\alpha([0, t])$ . Using the monotone class theorem for functions we get (69).

From [48, Corollary 5.2] we get that

$$\int P[\hat{\pi}] d\mathbf{Y} = \int P[\hat{\pi}] \circ dY,$$

and hence  $X^{\mathbf{Y}}$  satisfies (5). The result follows by uniqueness.  $\square$

## 5 The interacting particle system

In this section we prove well-posedness and convergence of the interacting particle system (13) in the case when  $Y$  is a path of bounded variation.

To compress the notation and have a slightly more general result we study the following mean-field system

$$dX_t^{i,N} = [b(X_t^{i,N}, \mu_t^N) + P(\mu_t^N)\dot{Y}_t + \Gamma(\mu_t^N)]dt + \sigma(X_t^{i,N}, \mu_t^N)dW_t^i, \quad X_t^{i,N}|_{t=0} = X_0^i, \quad i = 1, \dots, N. \quad (70)$$

The variable  $X_t^{i,N}$  is in the state space  $\mathbb{R}^D$ .  $(W^i)_{i \in \mathbb{N}}$  is a family of independent  $m$ -dimensional Brownian motions and  $(X_0^i)_{i \in \mathbb{N}}$  is a family of independent and identically distributed initial conditions with law  $\pi_0 \in \mathcal{P}(\mathbb{R}^D)$ . Moreover, assume that  $\dot{Y} : [0, T] \rightarrow \mathbb{R}^D$  is cadlag and bounded. We assume that the coefficients  $b$ ,  $\sigma$  and  $P$  satisfy Assumption 3, but notice that  $P$  only depends on the measure. For the coefficient  $\Gamma$  we assume the following

**Assumption 5.** Let  $\rho \geq 1$ , assume that there exists a constant  $C > 0$  such that

(i) (linear growth)  $\forall \pi \in \mathcal{P}_\rho(\mathbb{R}^D)$ ,

$$|\Gamma(\pi)| \leq C(1 + \overline{M}^\rho(\pi))^{\frac{2}{\rho}},$$

(ii) (Lipschitz continuity)  $\forall \pi, \nu \in \mathcal{P}_\rho(\mathbb{R}^D)$ ,

$$|\Gamma(\pi) - \Gamma(\nu)| \leq C(1 + \overline{M}^\rho(\pi)^2 + \overline{M}^\rho(\nu))^{\frac{1}{\rho}} W_\rho(\pi, \nu).$$

*Remark 5.1.* Notice that the growth is quadratic in the central moments of the measure. Also, the best Lipschitz constant we can hope for is the square of the central moment of one measure and the central moment of the other.

*Remark 5.2.* By doing the same substitution as in Section 4.5 and taking (9) for  $\Gamma$  we recover the interacting particle system (13).

**Lemma 5.3.** Let  $T > 0$  and  $\rho \geq 1$ . If the initial distribution  $\pi_0$  has finite  $\rho$ -moment, then equation (70) admits a pathwise unique strong solution on  $[0, T]$ . Moreover, there exists  $C = C(\rho, \pi_0, T) > 0$  such that

$$\frac{1}{N} \sum_{j=1}^N \mathbb{E} \left[ \sup_{t \in [0, T]} |\overline{X}_t^{j,N}|^\rho \right] \leq C, \quad \max_{i=1, \dots, N} \mathbb{E} \left[ \sup_{t \in [0, T]} |X_t^{i,N}|^\rho \right] \leq C(1 + \|\dot{Y}\|_\infty), \quad (71)$$

where  $\overline{X}^{i,N} = X^{i,N} - \frac{1}{N} \sum_{j=1}^N X^{j,N}$ , for  $i = 1, \dots, N$ .

*Proof.* Under Assumptions 3 and 5, the coefficients are locally Lipschitz and there is classically strong existence and pathwise uniqueness for the SDE (70) up to an explosion time  $\tau$ . We want to prove that  $\tau < T$ . Let us call the solution  $X_t^{(N)} = (X_t^{1,N}, \dots, X_t^{N,N})$ . Since the coefficients  $P$  and  $\Gamma$  do not depend on the state variable  $x$  we have the following identity for  $\overline{X}^{i,N}$ ,  $i = 1, \dots, N$ ,

$$d\overline{X}_t^{i,N} = (b(X_t^{i,N}, \mu_t^N) - \frac{1}{N} \sum_{j=1}^N b(X_t^{j,N}, \mu_t^N))dt + \sigma(X_t^{i,N}, \mu_t^N)dW_t^i - \frac{1}{N} \sum_{j=1}^N \sigma(X_t^{j,N}, \mu_t^N)dW_t^j.$$

Notice that

$$\overline{M}^\rho(\mu_t^N) = \mathbb{E}_{\mu_t^N} [|X - \mathbb{E}_{\mu_t^N}[X]|^\rho] = \frac{1}{N} \sum_{i=1}^N |\overline{X}_t^{i,N}|^\rho.$$

Using Assumption 3 we obtain that for every  $\rho \geq 2$  there exists a constant  $C > 0$  independent of  $\dot{Y}$  and  $N$  such that

$$\mathbb{E}[\sup_{s \leq t} |\overline{X}_s^{i,N}|^\rho] \leq \mathbb{E}[|\overline{X}_0^{i,N}|^\rho] + C \int_0^t (1 + \frac{1}{N} \sum_{j=1}^N \mathbb{E}[|\overline{X}_s^{j,N}|^\rho]) ds.$$

Taking the mean over  $i$  and using Gronwall's lemma we obtain the first estimate in (71). We can now estimate the moments of  $X^{i,N}$  as follows,

$$\mathbb{E} \left[ \sup_{t \in [0, T]} |X_t^{i,N}|^\rho \right] \leq C(1 + \|\dot{Y}\|_\infty) \int_0^T \mathbb{E} \left[ (1 + \overline{M}^\rho(\mu_s^N)) \right]^2 ds \leq (1 + \|\dot{Y}\|_\infty) C,$$

where  $C = C(\rho, \pi_0, T)$  can change from one inequality to the next. Since  $X^{(N)}$  has finite  $L_\omega^\rho$ -norm on the interval  $[0, T]$ , it means that the explosion time satisfies  $\tau > T$ .  $\square$

*Remark 5.4.* Notice that in Lemma 5.3 the choice of  $T > 0$  was arbitrary, which implies that the system of interacting particles is well posed on intervals of any length.

## 5.1 Propagation of chaos

For  $i = 1, \dots, N$ , we introduce the following McKean-Vlasov equation

$$dX_t^i = [b(X_t^i, \pi_t) + P(\pi_t)\dot{Y}_t + \Gamma(\pi_t)]dt + \sigma(X_t^i, \pi_t)dW_t^i, \quad \pi_t = \mathcal{L}(X_t^i), \quad X_t^i|_{t=0} = X_0^i \sim \pi_0. \quad (72)$$

By construction the random variables  $X^1, \dots, X^N$  are independent and identically distributed.

Equation (72) is well-posed because it corresponds to equation (61) driven by the *rough* path

$$\mathbf{Y}_{st} := (Y_{s,t}, \int_s^t Y_{s,r} \dot{Y}_r dr - \frac{1}{2}(t-s)I_{d \times d}),$$

where  $Y_t := \int_0^t \dot{Y}_r dr$ . Notice that, since  $\dot{Y}$  is cadlag and bounded, the path  $\mathbf{Y}$  is Lipschitz continuous which is much more regular than  $\alpha$ -Hölder with  $\alpha \in (\frac{1}{3}, \frac{1}{2}]$ .

Let  $\bar{X}^i := X^i - \mathbb{E}[X^i]$  and notice that, by definition  $\bar{M}^\rho(\pi_t) = \|\bar{X}_t^i\|_{L^\rho_\omega}^\rho$ . Using the a priori estimate in Lemma 4.7 and Assumptions 3 and 5 we can see that, for any  $\rho \geq 2$  we have an estimate on the central moments of  $\pi_t$ ,

$$\bar{M}^\rho(\pi_t) \leq \|\sup_{t \in [0, T]} \|\bar{X}_t^i\|_{L^\rho_\omega}^\rho \leq C[1 + \bar{M}^\rho(\pi_0)]^\rho.$$

From this one can easily recover the following estimate for the moments of  $\pi_t$ ,

$$M^\rho(\pi_t)^{\frac{1}{\rho}} \leq C(1 + \|\dot{Y}\|_\infty)(M^\rho(\pi_0) + \bar{M}^{2\rho}(\pi_t))^{\frac{1}{\rho}} \leq C(M^\rho(\pi_0) + \|\dot{Y}\|_\infty)(1 + \bar{M}^{2\rho}(\pi_0))^2, \quad (73)$$

where  $C = C(\rho, T)$ .

*Remark 5.5.* Notice that the  $\rho$ -moment of  $\pi_t$  is only bounded by the  $2\rho$ -central moment of  $\pi_0$  because of the quadratic growth of  $\Gamma$  from Assumption 5.

In the following we will hide the dependence on  $\pi_0$  in the constant  $C$  and only focus on the explicit dependence on  $\|\dot{Y}\|_\infty$ . We define the empirical measure associated with the independent particles  $X^i$  as  $\bar{\mu}_t^N := \frac{1}{N} \sum_{i=1}^N \delta_{X_t^i}$ . It follows from [46, Theorem 1] that for any  $\bar{\rho} > \rho$  there exists an explicit rate of convergence  $H(N) = H(N, \rho, \bar{\rho})$  such that  $H(N) \rightarrow 0$  as  $N \rightarrow \infty$  and

$$\mathbb{E}[W_{\rho, \mathbb{R}^D}(\bar{\mu}_t^N, \pi_t)^\rho] \leq \bar{M}^{\bar{\rho}}(\pi_t)^{\frac{\rho}{\bar{\rho}}} H(N) \leq C(1 + \|\dot{Y}\|_\infty) H(N), \quad (74)$$

where in the last inequality we used (73) and  $C$  depends on the moments of  $\pi_0$ .

Moreover, the rate is optimal and explicitly given as

$$H(N) = \begin{cases} N^{-\frac{1}{2}} + N^{1-\frac{\rho}{\bar{\rho}}} & \rho > \frac{D}{2}, \bar{\rho} \neq 2\rho \\ N^{-\frac{\rho}{D}} + N^{1-\frac{\rho}{\bar{\rho}}} & 1 \leq \rho < \frac{D}{2}, \bar{\rho} \neq \frac{D}{D-2}. \end{cases}$$

Notice that if we have enough moments the optimal rate of convergence for independent particles is  $H(N) = \frac{1}{\sqrt{N}}$ . Furthermore, we have the following estimate, because  $\mu^N$  and  $\bar{\mu}^N$  are both empirical measures,

$$W_{\rho, \mathbb{R}^D}(\mu_t^N, \bar{\mu}_t^N)^\rho \leq \frac{1}{N} \sum_{i=1}^N |X_t^{i,N} - X_t^i|^\rho,$$

so that we have from the triangular inequality

$$\sup_{t \in [0, T]} \mathbb{E}[W_{\rho, \mathbb{R}^D}(\mu_t^N, \pi_t)^\rho] \leq C(1 + \|\dot{Y}\|_\infty) H(N) + \sup_{t \in [0, T]} \frac{1}{N} \sum_{i=1}^N \mathbb{E}|X_t^{i,N} - X_t^i|^\rho, \quad (75)$$

where  $C$  depends on the  $2\bar{\rho}$ -moment of  $\pi_0$ , for  $\bar{\rho} > \rho$ . Now that we stated most of the preliminaries we can prove the following convergence result.

**Proposition 5.6.** *Let  $\bar{\rho} > \rho \geq 2$ . If  $X_0 \in L_{\omega}^{2\bar{\rho}}$ , there is a function  $J(N) = O(\log(N)^{-1})$  such that*

$$\max_{i=1, \dots, N} \mathbb{E}[\sup_{t \in [0, T]} |X_t^{i,N} - X_t^i|^\rho] \lesssim e^{\|\dot{Y}\|_\infty} J(N).$$

*Proof.* We call  $\bar{X}^{i,N} := X^{i,N} - \frac{1}{N} \sum_{j=1}^N X^{j,N}$ , for  $i = 1, \dots, N$  and recall that the central moment for the empirical measure is  $\bar{M}^\rho(\mu_t^N) = \frac{1}{N} \sum_{i=1}^N |\bar{X}_t^{i,N}|^\rho$ . For  $R > 0$ , we define the stopping time  $T_R := \inf\{t \geq 0 : \bar{M}^\rho(\mu_t^N) \geq R\}$ .

We set  $Z^i := X^{i,N} - X^i$  and using Assumptions 3 and 5 as well as inequality (75) and Lemma 4.7 we compute the following,

$$\begin{aligned} \max_{i=1,\dots,N} \mathbb{E}[|Z_t^i|^\rho 1_{\{T_R > t\}}] &\leq C \mathbb{E} \left[ \int_0^{t \wedge T_R} (1 + \overline{M}^{2\rho}(\pi_s) \|\dot{Y}\|_\infty + \overline{M}^\rho(\mu_s^N)) \left( |Z_s^i| + W_{\rho, \mathbb{R}^D}(\pi_s, \mu_s^N) \right)^\rho ds \right] \\ &\leq C(1 + \|\dot{Y}\|_\infty + R) \left[ \int_0^t \max_{i=1,\dots,N} \mathbb{E}[|Z_s^i|^\rho 1_{\{T_R \geq s\}}] ds + H(N) \right]. \end{aligned}$$

Using Gronwall's inequality we obtain

$$\max_{i=1,\dots,N} \mathbb{E}[|Z_t^i|^\rho 1_{\{T_R > t\}}] \leq C(1 + \|\dot{Y}\|_\infty) e^{C(1 + \|\dot{Y}\|_\infty)} R e^{CR} H(N).$$

Now we compute the following using Cauchy-Schwarz and Markov inequalities as well as (71),

$$\mathbb{E}[|Z_t^i|^\rho 1_{\{T_R \leq t\}}] \leq C \left( \mathbb{E}[|Z_t^i|^{2\rho}] \right)^{\frac{1}{2}} \mathbb{P}(T_R \leq t)^{\frac{1}{2}} \leq C \mathbb{P}(\sup_{s \leq t} \overline{M}^\rho(\mu_s^N) \geq R)^{\frac{1}{2}} \leq \frac{C}{R^{\frac{1}{2}}} \left( \frac{1}{N} \sum_{i=1}^N \mathbb{E} \left[ \sup_{t \in [0, T]} |\overline{X}_t^{i,N}|^\rho \right] \right)^{\frac{1}{2}} \leq \frac{C}{R^{\frac{1}{2}}}.$$

Notice that we used  $\mathbb{E}[|Z_t^i|^{2\rho}] \leq C(\mathbb{E}[|X_t^{i,N}|^{2\rho}] + \mathbb{E}[|X_t^i|^{2\rho}]) \leq C(1 + \|\dot{Y}\|_\infty)$ , hence we need from Remark 5.5 that the initial measure  $\pi_0$  has finite  $2\rho$  moments. We can put together the estimates to obtain

$$\max_{i=1,\dots,N} \mathbb{E}[|Z_t^i|^\rho] \leq \frac{C}{R^{\frac{1}{2}}} + C(1 + \|\dot{Y}\|_\infty + R) e^{C(1 + \|\dot{Y}\|_\infty)} e^{CR} H(N) \leq C e^{C(1 + \|\dot{Y}\|_\infty)} \left( \frac{1}{R^{\frac{1}{2}}} + e^{CR} H(N) \right),$$

where we changed the constants from one line to the next. Now choose  $R = R(N)$  such that

$$R(N) \rightarrow \infty, \quad e^{CR(N)} H(N) \rightarrow 0, \quad \text{as } N \rightarrow \infty.$$

Remember that  $H(N) \approx N^{-\gamma}$  with  $\gamma < \frac{1}{2}$  so that choosing  $R(N) \approx \log(N^{\overline{\gamma}})$  for some  $0 < \overline{\gamma} < \frac{\gamma}{C}$  we have a rate of convergence

$$J(N) \approx (\overline{\gamma} \log(N))^{-1} + N^{\overline{\gamma}C - \gamma} \approx \log(N)^{-1}.$$

□

*Remark 5.7.* Notice that the rate of convergence is far from the optimal  $\frac{1}{\sqrt{N}}$  of a sample of independent and identically distributed random variables. This is due to the non-local Lipschitz condition on  $\Gamma$  in Assumption 5.

Finally, we can put together Corollary 4.16 and Proposition 5.6.

**Theorem 5.8.** *Let  $\delta > 0$  and  $\mathbf{Y}^\delta$  be a bounded differentiable approximation of a geometric rough path  $\mathbf{Y} = (Y, \mathbb{Y}) \in \mathcal{C}^{\overline{\alpha}}$ . Let  $\mu^{N,\delta}$  be the empirical measure of the system of mean-field particles (70) with input  $\mathbf{Y}^\delta$ . Moreover, let  $\pi$  be the law of the solution to equation (61) driven by  $(Y, \mathbb{Y} - \frac{1}{2}(t-s)\text{Id}) \in \mathcal{C}^{\overline{\alpha}}$ .*

*Then there exists  $\rho > 0$  and a sequence  $\delta(N)$  such that  $\delta(N) \rightarrow 0$  and*

$$\sup_{t \in [0, T]} \mathbb{E}[W_{\rho, \mathbb{R}^D}(\mu_t^{N,\delta(N)}, \pi_t)^\rho] \rightarrow 0, \quad \text{as } N \rightarrow \infty.$$

*Proof.* Let  $\pi^\delta$  be the law of a solution to (72). By the triangular inequality, Proposition 5.6 and Corollary 4.16

$$\begin{aligned} \sup_{t \in [0, T]} \mathbb{E}[W_{\rho, \mathbb{R}^D}(\mu_t^{N,\delta(N)}, \pi_t)^\rho] &\leq \sup_{t \in [0, T]} \mathbb{E}[W_{\rho, \mathbb{R}^D}(\mu_t^{N,\delta(N)}, \pi_t^\delta)^\rho] + W_{\rho, C([0, T], \mathbb{R}^D)}(\pi^\delta, \pi) \\ &\lesssim e^{\|\dot{Y}^\delta\|_\infty} J(N) + \rho_{\overline{\alpha}}(\mathbf{Y}^\delta, \mathbf{Y}). \end{aligned}$$

Choosing  $\delta(N)$  such that  $\|\dot{Y}^{\delta(N)}\|_\infty = o(\log(J(N)^{-1})) = o(\log(\log(N)))$  we have that the right hand side vanishes as  $N \rightarrow \infty$ . □

## 6 The numerical scheme, construction of the lift and examples

In this section we derive the RP-EnKF (rough path ensemble Kalman filter) numerical scheme alluded to in the introduction (see equation (14)), discuss the construction of appropriate rough path lifts, and provide details concerning the implementation. Furthermore, we demonstrate its effectiveness in the context of misspecified and multiscale models by means of a few examples in the context of parameter estimation. This setting provides a convenient testbed for the scenario where the model and observation noises are correlated, and we expect our conclusions regarding (non-)robustness to be relevant more generally.

**Discretising in time.** A natural scheme for approximating the dynamics of the interacting particle system (13) is given by

$$X_{k+1}^i = X_k^i + f(X_k^i)\Delta t + G^{1/2}\sqrt{\Delta t}\xi_k^i + \hat{P}_k \left( \Delta Y_k - (h(X_k^i)\Delta t + U\sqrt{\Delta t}\xi_k^i + R^{1/2}\sqrt{\Delta t}\eta_k^i) \right) \quad (76a)$$

$$+ \widehat{\text{Cov}}_k(x, Dh)\hat{P}_k\Delta\mathbb{Y}_k + \hat{\Gamma}_k\Delta t, \quad (76b)$$

where  $\Delta t > 0$  is the step size, and  $(\xi_n^i)$  and  $(\eta_n^i)$  denote independent zero mean Gaussian random variables with unit variance of dimensions  $D$  and  $d$ , respectively. In the above display,  $\hat{P}_k := \widehat{\text{Cov}}_k(x, h)C^{-1} + B$  and  $\widehat{\text{Cov}}_k(x, Dh)$  refer to the standard unbiased empirical estimators of the covariance, that is,

$$\widehat{\text{Cov}}_k(x, h) = \frac{1}{N-1} \sum_{i=1}^N (X_k^i - \bar{X}_k) \otimes h(X_k^i) \in \mathbb{R}^{D \times d}, \quad (77a)$$

$$\widehat{\text{Cov}}_k(x, Dh) = \frac{1}{N-1} \sum_{i=1}^N (X_k^i - \bar{X}_k) \otimes Dh(X_k^i) \in \mathbb{R}^{D \times d \times D}, \quad (77b)$$

where

$$\bar{X}_k = \frac{1}{N} \sum_{i=1}^N X_k^i \quad (78)$$

refers to the empirical mean. With (77) in place, the standard empirical estimator for  $\Gamma$  as defined in (9) is given by

$$\hat{\Gamma}_{k,\gamma} = -\frac{1}{2} \text{Trace} \left( \widehat{\text{Cov}}_{k,\gamma}(x, Dh)\hat{P}_k \right), \quad (79)$$

where  $\hat{\Gamma}_{k,\gamma}$  denotes the  $\gamma^{\text{th}}$  component of  $\hat{\Gamma}_k \in \mathbb{R}^D$ , and using a similar convention for  $\widehat{\text{Cov}}_{k,\gamma}(x, Dh)$ . the precise meaning of the first term in equation (76b) is

$$(\widehat{\text{Cov}}_k(x, Dh)\hat{P}_k\Delta\mathbb{Y}_k)_\gamma = \sum_{j,q=1}^d \sum_{r=1}^D \widehat{\text{Cov}}_k(x, Dh)_{\gamma,j,r} (\hat{P}_k)_{r,q} (\Delta\mathbb{Y}_k)_{q,j} \quad \gamma = 1, \dots, D,$$

and analogously in equation (79) with  $\Delta\mathbb{Y}_k$  replaced by the  $d$ -dimensional identity matrix.

*Remark 6.1* (Gubinelli derivative). The first term in (76b) is modelled after the Gubinelli derivative  $P(\hat{\pi}_s)'$  in (12). We would like to stress that a standard time discretisation of the interacting particle system (13) according to Davie [31] would involve further contributions accounting for correlations between the particles as well as for cross terms induced by the joint lift  $(Y, W, V) \mapsto ((Y, W, V), (\mathbb{Y}, \mathbb{W}, \mathbb{V}))$ . At least formally, these additional terms vanish in the limit as  $N \rightarrow \infty$ , and our numerical experiments have not shown noticeable benefits of including them.

**Constructing the lift  $\mathbb{Y}$ .** In order to implement the RP-EnKF scheme defined in (76), we need to posit the discrete-time second order increments  $\Delta\mathbb{Y}_k \in \mathbb{R}^{d \times d}$ , given discrete-time samples  $y_0, y_1, \dots, y_n \in \mathbb{R}^d$  from  $(Y_t)_{0 \leq t \leq T}$ . In what follows we will denote the piecewise-linear interpolation of  $y_0, \dots, y_n$  by  $(\tilde{Y}_t)_{0 \leq t \leq T}$  and consider the decomposition of  $\Delta\mathbb{Y}_k$  into symmetric and skew-symmetric parts,

$$\Delta\mathbb{Y}_k = \Delta\mathbb{Y}_k^{\text{sym}} + \Delta\mathbb{Y}_k^{\text{skew}}, \quad (80)$$

where

$$\Delta\mathbb{Y}_k^{\text{sym}} := \text{sym}(\Delta Y_k) := \frac{1}{2} (\Delta\mathbb{Y}_k + \Delta\mathbb{Y}_k^\top), \quad \Delta\mathbb{Y}_k^{\text{skew}} := \text{skew}(\Delta Y_k) := \frac{1}{2} (\Delta\mathbb{Y}_k - \Delta\mathbb{Y}_k^\top). \quad (81)$$

For the symmetric part, we set

$$\Delta\mathbb{Y}_k^{\text{sym}} = \text{sym} \left( \int_{t_k}^{t_{k+1}} \tilde{Y}_{t_k, r} \otimes d\tilde{Y}_r \right) = \frac{1}{2} (y_{k+1} - y_k) \otimes (y_{k+1} - y_k), \quad (82)$$

maintaining structural similarities to the defining algebraic relation of weakly geometric rough paths [48, Section 2.2]. Defining the skew-symmetric part is more challenging since<sup>3</sup>

$$\text{skew} \left( \int_{t_k}^{t_{k+1}} \tilde{Y}_{t_k, r} \otimes d\tilde{Y}_r \right) = 0, \quad (83)$$

indicating that information on the enclosed area between two neighbouring points is inevitably lost by the discretisation. In some scenarios, sensible values for  $\Delta\mathbb{Y}_k^{\text{skew}}$  may be dictated by a specific argument related to

<sup>3</sup>This relation expresses the fact that the area enclosed by a straight line with itself is zero.

the particular nature of  $(Y_t)_{0 \leq t \leq T}$ , see Section 6.1. In other circumstances, setting  $\Delta \mathbb{Y}_k^{\text{skew}} = 0$  might provide a reasonable approximation, see Section 6.3. Failing that, our experiments show promising results based on the following procedure (for other approaches towards estimating Lévy areas see [9, 44, 71]):

For a specific time-lag  $\tau \in \mathbb{N}_{\geq 1}$ , consider the subsampled sequence  $y_0, y_\tau, y_{2\tau}, \dots$ , as well as the associated piecewise linear interpolation  $(\tilde{Y}_t^\tau)_{0 \leq t \leq T}$ . The time-lag  $\tau$  shall be chosen in such a way that the corresponding area paths

$$t \mapsto \text{skew} \left( \int_0^t \tilde{Y}_{0,r} \otimes d\tilde{Y}_r \right) =: \tilde{\mathbb{Y}}_{0,t}^{\text{skew}} \quad \text{and} \quad t \mapsto \text{skew} \left( \int_0^t \tilde{Y}_{0,r}^\tau \otimes d\tilde{Y}_r^\tau \right) =: \tilde{\mathbb{Y}}_{0,t}^{\text{skew},\tau} \quad (84)$$

are ‘as distinct as possible’, while maintaining  $(\tilde{Y}_t)_{0 \leq t \leq T} \approx (\tilde{Y}_t^\tau)_{0 \leq t \leq T}$ . These comparisons can be made in supremum norm, for instance. We then set

$$\Delta \mathbb{Y}_k^{\text{skew}} = (\tilde{\mathbb{Y}}_{0,t_{k+1}}^{\text{skew},\tau} - \tilde{\mathbb{Y}}_{0,t_k}^{\text{skew},\tau}) - (\tilde{\mathbb{Y}}_{0,t_{k+1}}^{\text{skew}} - \tilde{\mathbb{Y}}_{0,t_k}^{\text{skew}}). \quad (85)$$

Equation (85) compares the area contributions associated to the original interpolation  $\tilde{Y}$  and the subsampled interpolation  $\tilde{Y}^\tau$ . The requirement  $(\tilde{Y}_t)_{0 \leq t \leq T} \approx (\tilde{Y}_t^\tau)_{0 \leq t \leq T}$  is meant to ensure that  $\tilde{Y}$  and  $\tilde{Y}^\tau$  mainly differ at the second-order level  $\mathbb{Y}$ ; visually, the subsampling operation may be understood as ‘straightening out’  $\tilde{Y}$  and measuring the area accumulated thereby.

*Remark 6.2* (Relationship to subsampling in multiscale parameter estimation). As mentioned in Section 2.3, ideas related to subsampling have been considered extensively in the context of multiscale parameter estimation (without observational noise), see, for instance, [1, 5, 6, 51, 52, 61, 63, 64, 83, 88]. The method proposed in this section is different in that we use the subsampled paths in order to estimate the Lévy area, but otherwise input the original data path into the RP-EnKF dynamics (14). Moreover, the motivations are distinct: While subsampling in the aforementioned works is used in order to eliminate small-scale fluctuations, our method is specifically designed to estimate Lévy area terms. We leave an exploration of the connection between both methods for future work.

## 6.1 Physical Brownian motion in a magnetic field

In a first example, we consider a parameter estimation problem where the dynamics of interest is driven by a physical Brownian motion subject to a magnetic field. More precisely, physical Brownian motion  $(W_t^\varepsilon)_{t \geq 0}$  is defined in terms of the unique strong solution to the following system of SDEs,

$$dW_t^\varepsilon = \frac{1}{\varepsilon} M^{-1} P_t^\varepsilon dt, \quad W_0^\varepsilon = 0, \quad (86a)$$

$$dP_t^\varepsilon = -\frac{1}{\varepsilon} M P_t^\varepsilon dt + dW_t^0, \quad P_0^\varepsilon = 0, \quad (86b)$$

where  $W_t^\varepsilon, P_t^\varepsilon \in \mathbb{R}^2$ ,  $(W_t^0)_{t \geq 0}$  is a standard (mathematical) two-dimensional Brownian motion, and  $\varepsilon \ll 1$  is a small parameter, the limit  $\varepsilon \rightarrow 0$  corresponding to the regime of negligible particle mass. Furthermore, the matrix  $M$  is given by

$$M = \begin{pmatrix} 1 & \gamma \\ -\gamma & 1 \end{pmatrix}, \quad (87)$$

with  $\gamma \in \mathbb{R}$  being a real-valued parameter associated to the strength of the magnetic field. For fixed  $\alpha \in (1/3, 1/2)$  and  $T > 0$ , it is known that  $(W^\varepsilon, \mathbb{W}^\varepsilon) \rightarrow (W^0, \mathbb{W}^{\text{phys}}(\gamma))$  in  $\mathcal{C}^\alpha([0, T], \mathbb{R}^d)$  and  $L^1$  as  $\varepsilon \rightarrow 0$ , where

$$\mathbb{W}_{s,t}^\varepsilon = \int_s^t W_{s,r}^\varepsilon \otimes dW_r^\varepsilon \quad (88)$$

denotes the canonical lift, and

$$\mathbb{W}_{s,t}^{\text{phys}}(\gamma) = \int_s^t W_{s,r}^0 \otimes \odot dW_r^0 + (t-s)D, \quad (89)$$


with area correction

$$D = \frac{1}{2} \begin{pmatrix} 0 & \gamma \\ -\gamma & 0 \end{pmatrix}, \quad (90)$$

see [47] and [48, Section 3.4]. The setting is reminiscent of the passage between underdamped and overdamped Langevin dynamics, see [87, Section 6.5.1] and [88, Section 2.2]. Similarly to [37, Section 8.2], we consider the problem of estimating the parameter  $\theta \in \mathbb{R}$  in

$$dZ_t^\varepsilon = \theta f(Z_t^\varepsilon) dt + dW_t^\varepsilon, \quad Z_0^\varepsilon = 0, \quad (91)$$

Figure 1:  $t \mapsto (Z_t^\varepsilon)_{0 \leq t \leq T}$  and  $t \mapsto \int_0^t Z_s^\varepsilon \otimes dZ_s^\varepsilon$  for mathematical and physical Brownian motion.



given noisy observations of the path  $(Z_t^\varepsilon)_{0 \leq t \leq T}$ , that is, given a path  $(Y_t^\varepsilon)_{0 \leq t \leq T}$  of the solution to


$$dY_t^\varepsilon = dZ_t^\varepsilon + R^{1/2} dV_t, \quad Y_0 = 0, \quad (92)$$

see Section 2.3. In (91) and (92), we allow for both  $\varepsilon = 0$  and  $\varepsilon > 0$ , that is, we consider the dynamics driven by both mathematical and physical Brownian motion. Note that in the noiseless case  $R = 0$  our setting coincides with the one discussed in [37], see also Appendix B. Standard arguments show that both  $(Z^\varepsilon, \mathbb{Z}^\varepsilon)$  and  $(Y^\varepsilon, \mathbb{Y}^\varepsilon)$  converge in  $\mathcal{C}^\alpha([0, T], \mathbb{R}^d)$  with a nontrivial area correction akin to (89), where in the latter case,  $\mathbb{Y}^\varepsilon$  refers to the Stratonovich iterated integrals. As an illustration, we plot sample paths of  $t \mapsto Z_t^\varepsilon$  and an off-diagonal component of  $t \mapsto \int_0^t Z_s^\varepsilon \otimes dZ_s^\varepsilon$  in Figure 1, comparing the cases  $\varepsilon = 0$  and  $\varepsilon = 10^{-2}$ , for the same realisation of (a discretised version of)  $(W_t^0)_{0 \leq t \leq T}$ . Crucially, the convergence towards a nontrivially lifted path as expressed in (89) manifests itself in the offset between the paths in Figure (1b). Throughout this section, we choose a fine time step of  $\Delta t = 10^{-4}$  for all the involved approximations,  $\theta_{\text{true}} = 1/2$  for the parameter to be recovered,  $\gamma = -2.0$  for the strength of the magnetic field,  $R = 0.1$  for the variance of the observation noise, and  $f(z_1, z_2) = (z_1 - z_2, z_1 + z_2)^\top$  for the drift in (91).

To test the robustness of the EnKF scheme (17) and the RP-EnKF scheme (14), we generate data according to (91) and (92) for both  $\varepsilon = 0$  (mathematical Brownian motion) and  $\varepsilon = 10^{-2}$  (physical Brownian motion). We would like to stress that the filtering methodology (expressed in terms of the schemes (17) and (14)) is however based on the model (1) and therefore tailored to the case  $\varepsilon = 0$ . In Figure 2, we show the empirical mean of the  $\hat{\theta}$ -components for the output of the EnKF-dynamics (17), considering both mathematical and physical Brownian motion as drivers in (91). Evidently, the EnKF is not robust, in the sense that it fails to recover the true parameter  $\theta_{\text{true}}$  in the case when  $\varepsilon = 10^{-2}$ , see Figure (2b).

We proceed by showing the corresponding results for the RP-EnKF scheme defined by (76) in Figure 3, demonstrating the robustness promised by Theorem 4.17. For the required (discrete-time) lift  $\Delta \mathbb{Y}$ , we use the construction detailed in Section 6. More precisely, in the case of physical Brownian motion, we set the time-lag to  $\tau = 700$ . In the case of mathematical Brownian motion, we set the skew-symmetric part in (80) to zero,  $\Delta \mathbb{Y}_k^{\text{skew}} = 0$ , corresponding to the choice  $\tau = 1$ . These choices have been made on the basis of the subsampled area processes  $t \mapsto \text{skew} \left( \int_0^t \tilde{Y}_s^{\varepsilon, \tau} \otimes d\tilde{Y}_s^{\varepsilon, \tau} \right)$  depicted in Figure 4. More precisely, Figures (4a) and (4c) show the dependence  $t \mapsto \text{skew} \left( \int_0^t \tilde{Y}_s^{\varepsilon, \tau} \otimes d\tilde{Y}_s^{\varepsilon, \tau} \right)$  with different values of the time-lag  $\tau$ , for physical Brownian motion ( $\varepsilon = 10^{-2}$ , Figure (4a)) and mathematical Brownian motion ( $\varepsilon = 0$ , Figure (4c)). While the subsampling only minimally effects the area process associated to the process driven by mathematical Brownian motion (Figure 4c), we observe a systematic shift in the case of physical Brownian motion (Figure (4c)), revealing the latent multiscale structure. The difference between the original and subsampled area processes for physical Brownian motion is shown in Figure (4b). As the time-lag  $\tau$  increases, said difference approaches the theoretically expected area correction implied by (89)-(90). The fact that the difference between the original and the subsampled area process reaches a plateau at around  $\tau = 700$  is illustrated in Figure (4e), as opposed to the difference between the original and the subsampled paths, see Figure (4d). Consequently, the choice  $\tau = 700$  strikes a balance between separating the original and subsampled area processes as much as possible while maintaining similarity

Figure 2: Output (empirical mean of the  $\hat{\theta}$ -components) of the EnKF-scheme (17), for data obtained from the dynamics (91) driven by either mathematical or physical Brownian motion. The blue line indicates the true value  $\theta_{\text{true}} = \frac{1}{2}$ . To suppress sampling error, we plot averaged results computed from 5 independent runs.



between the original and subsampled paths (as suggested by the discussion motivating (85)).

## 6.2 Fast chaotic dynamics – Lorenz-63

In this example, we consider the rescaled Lorenz ordinary differential equations [94] for  $L_t^\varepsilon = (L_t^{(1),\varepsilon}, L_t^{(2),\varepsilon}, L_t^{(3),\varepsilon}) \in \mathbb{R}^3$ ,

$$\dot{L}_t^{(1),\varepsilon} = \frac{\sigma}{\varepsilon^2} (L_t^{(2),\varepsilon} - L_t^{(1),\varepsilon}), \quad L_0^{(1),\varepsilon} = l_0^{(1)}, \quad (93a)$$

$$\dot{L}_t^{(2),\varepsilon} = \frac{1}{\varepsilon^2} (\rho L_t^{(1),\varepsilon} - L_t^{(2),\varepsilon} - L_t^{(1),\varepsilon} L_t^{(3),\varepsilon}), \quad L_0^{(2),\varepsilon} = l_0^{(2)}, \quad (93b)$$

$$\dot{L}_t^{(3),\varepsilon} = \frac{1}{\varepsilon^2} (L_t^{(1),\varepsilon} L_t^{(2),\varepsilon} - \beta L_t^{(3),\varepsilon}), \quad L_0^{(3),\varepsilon} = l_0^{(3)}, \quad (93c)$$

with the standard parameters  $\sigma = 10$ ,  $\rho = 28$  and  $\beta = \frac{8}{3}$  as an example of fast chaotic dynamics approximating Brownian noise (with nontrivial area correction). We would like to stress that the phenomena described in this section can be considered generic across a wide range of fast chaotic deterministic dynamical systems, and refer the reader to [20] for an overview.

With  $\varepsilon = 1$ , the system (93) has originally been proposed as a simplified model for atmospheric convection [75], and can serve as a prototype for the study of chaotic ODEs. As is well known, (93) possesses a ‘strange’ chaotic attractor  $\Lambda$  equipped with a unique SRB (Sinai-Ruelle-Bowen) measure<sup>4</sup>  $\mu$ , see [104]. For  $\varepsilon \ll 1$ , a random initial condition  $L_0^\varepsilon \in \mathbb{R}^3$  and after appropriate centering and rescaling, the solution  $(L_t)_{t \geq 0}$  is well approximated by a Brownian motion with a nontrivial area correction in the sense of rough paths:

For Hölder-continuous observables  $v : \mathbb{R}^3 \rightarrow \mathbb{R}^m$  that are  $\mu$ -centred (that is,  $\int_{\mathbb{R}^3} v \, d\mu = 0$ ), a functional CLT (or *weak invariance principle*) holds for

$$W_t^\varepsilon := \frac{1}{\varepsilon} \int_0^t v(L_s^\varepsilon) \, ds, \quad (94)$$


assuming that  $L_0^\varepsilon$  is initialised randomly according to  $\mu$ . More precisely, there exists a Brownian motion  $W$  (possibly on an extended probability space) with appropriate covariance  $\Sigma \in \mathbb{R}^{m \times m}$  such that  $W^\varepsilon \rightarrow W$  weakly in  $C([0, T], \mathbb{R}^m)$  as  $\varepsilon \rightarrow 0$ , see [58, Theorem 1.5]<sup>5</sup>. Moreover, it was shown in [11, Theorem 1.6] that a so-called *iterated weak invariance principle* holds for the iterated integrals

$$\mathbb{W}_t^\varepsilon = \int_0^t W_s^\varepsilon \otimes dW_s^\varepsilon, \quad (95)$$

<sup>4</sup>The notion of SRB measures provides a suitable generalisation of ergodic measures.

<sup>5</sup>In fact, the convergence takes place  $\mu$ -almost surely, see [58, Theorem 1.1]. The covariance  $\Sigma$  is given in terms of suitable long-time ergodic averages or (under certain conditions) Green-Kubo formulae, see [20, 86].

Figure 3: Output (empirical mean of the  $\hat{\theta}$ -components) of the RP-EnKF-scheme (14), for data obtained from the dynamics (91) driven by either mathematical or physical Brownian motion. The blue line indicates the true value  $\theta_{\text{true}} = \frac{1}{2}$ . To suppress sampling error, we plot averaged results computed from 5 independent runs.



that is  $(W^\varepsilon, \mathbb{W}^\varepsilon) \rightarrow (W, \mathbb{W})$  weakly in  $C([0, T], \mathbb{R}^m \times \mathbb{R}^{m \times m})$ , where

$$\mathbb{W}_t = \int_0^t W_s \otimes \circ dW_s + Dt, \quad (96)$$

with area correction  $D \in \mathbb{R}^{m \times m}_{\text{skew}}$ . We refer to [20, Theorem 4.4] for the corresponding statement in  $p$ -variation rough path topology.

In what follows, we consider  $(Z_t^\varepsilon)_{t \geq 0}$  to be driven by  $(L_t^{(1:2), \varepsilon})_{t \geq 0} := (L_t^{(1), \varepsilon}, L_t^{(2), \varepsilon})_{t \geq 0}$ ,

$$dZ_t^\varepsilon = \theta f(Z_t^\varepsilon) dt + \frac{\lambda}{\varepsilon} dL_t^{(1:2), \varepsilon}, \quad (97)$$

with the parameter  $\theta \in \mathbb{R}$  to be inferred from noisy observations


$$dY_t^\varepsilon = dZ_t^\varepsilon + R^{1/2} dV_t, \quad (98)$$

and  $\lambda > 0$  mediating the strength of the chaotic perturbation. Since  $\int_{\mathbb{R}^d} (L^{(1)}, L^{(2)})^\top d\mu = 0$  according to [86, Section 11.7.2], we expect (97) to be well approximated by


$$dZ_t^0 = \theta f(Z_t^0) dt + G^{1/2} dW_t, \quad (99)$$

in the regime  $\varepsilon \ll 1$ , with standard two-dimensional Brownian motion  $(W_t)_{t \geq 0}$  and appropriate covariance  $G \in \mathbb{R}^{2 \times 2}$ . Replacing (97) by (99) is often a desirable simplification both computationally and conceptually, see, for instance, [53], [86, Section 11.7.2], and [99, Section 3.1]. To apply our methodology, we need to presuppose  $G$ ; estimates can be obtained using the approaches suggested in [53, Example 6.2] or [63], for instance. Here we use the value  $G_{11}^{1/2} = 0.13$  reported in [63, Section 3.2.6] for  $\lambda = \frac{2}{45}$  and note that  $W^1 = W^2$  almost surely by a short calculation using (93). To test the RP-EnKF, we simulate data according to (93), (97) and (98) using an Euler-Maruyama discretisation with time step  $\Delta t = 10^{-5}$ , an observation noise level of  $R = 0.01$ , a parameter value of  $\theta_{\text{true}} = 0.5$  to be recovered. Furthermore, we set  $\varepsilon = 0.05$  and  $f(z_1, z_2) = (z_1 - z_2, z_1 + z_2)^\top$ . The RP-EnKF is set up according to the model (99) with observations (98), using the scheme (14). As an illustration for the challenges that are posed by the attempt to incorporate data from (97)-(98) into a model of the form (99), we plot the output of the EnKF scheme (17) in Figure (5a), noting that it fails to recover the correct parameter value  $\theta_{\text{true}}$ . Figures (5b) and (5c) show the output of the RP-EnKF scheme (14), with  $\tau = 1$  (implying  $\Delta Y_k^{\text{skew}} = 0$ ) and  $\tau = 500$ , respectively. We see that the area correction obtained through the subsampling procedure is necessary to obtain satisfactory numerical results. The time-lag  $\tau = 500$  has been determined in the same way as in Section 6.1; in particular, plots for the subsampled area processes are qualitatively similar to Figures 4 and 4b but are omitted here for the sake of brevity.

Figure 4: Illustration of the subsampling procedure from Section 6 for constructing the discrete-time lift  $\Delta \mathbb{Y}$




(a) Area processes for physical Brownian motion ( $\varepsilon = 10^{-2}$ ). (b) Differences of subsampled and original area processes ( $\varepsilon = 10^{-2}$ ). (c) Area processes for mathematical Brownian motion ( $\varepsilon = 0$ ).



(d)  $L^2$ -discrepancy between original and subsampled path. (e)  $L^2$ -discrepancy between original and subsampled area process.


- (4a): Area processes  $t \mapsto \text{skew} \left( \int_0^t \tilde{Y}_s^{\varepsilon, \tau} \otimes d\tilde{Y}_s^{\varepsilon, \tau} \right)$  (off-diagonal component) for physical Brownian motion ( $\varepsilon = 10^{-2}$ ) and time-lags  $\tau = 100$ (green),  $\dots$ , 800 (yellow). The area processes associated to the path without subsampling is plotted in black.
- (4b): Differences of original and subsampled area processes in the case of physical Brownian motion ( $\varepsilon = 10^{-2}$ ), that is  $t \mapsto \int_0^t \left( \tilde{Y}_s^{\varepsilon, \tau} \otimes d\tilde{Y}_s^{\varepsilon, \tau} - Y_s^{\varepsilon} \otimes dY_s^{\varepsilon} \right)$ , in the spirit of (85). The dashed line represents the theoretically expected area correction according to (89).
- (4c): Area processes associated to mathematical Brownian motion ( $\varepsilon = 0$ ), same colour scheme as for Figure (4a).
- (4d):  $L^2$ -discrepancy between the original path  $(Y_t)_{0 \leq t \leq T}$  and the subsampled path  $(\tilde{Y}_t^{\tau})_{0 \leq t \leq T}$  as a function of the time-lag  $\tau$  for physical Brownian motion ( $\varepsilon = 10^{-2}$ ).
- (4e):  $L^2$ -discrepancy between the original area process  $\text{skew} \left( \int_0^t Y_s \otimes dY_s \right)_{0 \leq t \leq T}$  and the subsample area process  $\text{skew} \left( \int_0^t \tilde{Y}_s^{\tau} \otimes d\tilde{Y}_s^{\tau} \right)_{0 \leq t \leq T}$  as a function of the time-lag  $\tau$  for physical Brownian motion ( $\varepsilon = 10^{-2}$ ).

Figure 5: Output (empirical mean of the  $\hat{\theta}$ -components) EnKF and RP-EnKF (with and without area correction), for data obtained from the dynamics (97) perturbed by fast chaotic noise (Lorenz-63). The blue line indicates the true value of  $\theta$ . To suppress sampling error, average results are plotted computed from 5 independent runs.



(a) EnKF according to (17). (b) RP-EnKF without area correction. (c) RP-EnKF with area correction.

Figure 6: Output (empirical mean of the  $\hat{\theta}$ -components) of the EnKF-scheme (17), for data obtained from the reduced model (101) (left) and the multiscale model (100) (right). The blue line indicates the true value  $\theta_{\text{true}} = 1$ . To suppress sampling error, we plot averaged results computed from 5 independent runs.



### 6.3 Homogenisation in a two-scale potential

In our final example, we consider the motion of a Brownian particle in a (rugged) two-scale potential,

$$dZ_t^\varepsilon = -\theta \nabla V(Z_t^\varepsilon) dt - \frac{1}{\varepsilon} \nabla p\left(\frac{Z_t^\varepsilon}{\varepsilon}\right) dt + \sqrt{2\sigma} dW_t, \quad (100)$$

where  $p = (p_1, p_2) : \mathbb{R}^2 \rightarrow \mathbb{R}$  is an  $L$ -periodic function in both directions, that is,  $p_i(x + L) = p_i(x)$ , for all  $x \in \mathbb{R}$  and  $i = 1, 2$ , modelling small-scale fluctuations around the potential  $V$ . It is well known that for  $T > 0$ , the law of the solution  $(Z_t^\varepsilon)_{t \geq 0}$  converges weakly in  $C([0, T]; \mathbb{R}^d)$  to the law associated to

$$dZ_t = -\theta \mathcal{K} \nabla V(Z_t, \theta) dt + \sqrt{2\sigma \mathcal{K}} dW_t, \quad (101)$$

where  $\mathcal{K} = \text{diag}(L^2/(C_1 \hat{C}_1), L^2/(C_2 \hat{C}_2))$ , with


$$C_i = \int_0^L e^{-\frac{p_i(y)}{\sigma}} dy, \quad \hat{C}_i = \int_0^L e^{\frac{p_i(y)}{\sigma}} dy, \quad (102)$$

see [89] and [86, Chapter 11]. Similar results in a rough-path context can be found in [69]. The homogenised dynamics (101) encapsulates the rugged landscape described by  $p$  in the diffusion-mass matrix  $\mathcal{K}$ . Like in the previous experiments, we consider the task of estimating the parameter  $\theta$  from noisy observations of (100), using the EnKF and RP-EnKF based on (101). We choose  $V(z) = \frac{1}{2}|z|^2$ ,  $p_1(x) = \cos(x)$ ,  $p_2(x) = \frac{1}{2}\cos(x)$  and  $\sigma = 1$ . Data from the two-scale dynamics (100) is simulated for  $\varepsilon = 10^{-2}$ ,  $\theta_{\text{true}} = 1$  using an Euler-Maruyama discretisation with time step  $\Delta t = 10^{-4}$ . With the same values for  $\theta_{\text{true}}$ ,  $\Delta t$  and  $\sigma$ , we simulate data from the reduced model (101), where  $\mathcal{K} \approx \text{diag}(0.62386, 0.884176)$  has been obtained by numerical integration in (102). Both (100) and (101) are perturbed by noise according to (98) with  $R = 10^{-2}$  and, as in Sections 6.1 and 6.2, the resulting observation paths are used in the EnKF- and RP-EnKF schemes (see equations (17) and (14)) to estimate  $\theta_{\text{true}}$ . We display the results for the means of  $\hat{\theta}$  over time using the EnKF and the RP-EnKF in Figures 6 and 7, respectively.

Clearly, the RP-EnKF deals adequately with multiscale data, while the EnKF fails to recover the true parameter  $\theta_{\text{true}}$  in this setting. We would like to stress that the RP-EnKF scheme has been implemented without area correction, that is imposing  $\mathbb{Y}_k = \mathbb{Y}_k^{\text{sym}}$  as defined in (82). The choice  $\mathbb{Y}_k^{\text{skew}} = 0$  is motivated by a plot analogous and qualitatively similar to Figure 4c (omitted due to space considerations), showing that subsampling does not indicate substantial Lévy area terms (intuitively, the dynamics (100) does not contain significant ‘rotational’ contributions in the regime  $\varepsilon \rightarrow 0$ ).

**Acknowledgements.** This research has been partially funded by Deutsche Forschungsgemeinschaft (DFG) through the grant CRC 1114 ‘Scaling Cascades in Complex Systems’ (project A02, project number 235221301). We would like to thank R. Chew, A. Djurdjevac, G. Hastermann, R. Klein and S. Reich for very stimulating discussions. N.N. would like to thank M. Engel for organising a seminar on homogenisation at Imperial College London in 2016 leading to Section 6.2, as well as for sharing his insight into the topic.

Figure 7: Output (empirical mean of the  $\hat{\theta}$ -components) of the RP-EnKF-scheme (14), for data obtained from the reduced model (101) (left) and the multiscale model (100) (right). No area correction is used, that is,  $\mathbb{Y}_k^{\text{skew}} = 0$ . The blue line indicates the true value  $\theta_{\text{true}} = 1$ . To suppress sampling error, we plot averaged results computed from 5 independent runs.



## A From the filter to the McKean-Vlasov

In this section we prove the formal connection between the filtering problem and the McKean-Vlasov equation. In order to prove Proposition 2.1, we need a few preparations. Let us define

$$M_t = \int_0^t h(X_s) \cdot C^{-1} dY_s,$$

and the likelihood

$$l_t = \exp \left( M_t - \frac{1}{2} \langle M \rangle_t \right) = \exp \left( \int_0^t h(X_s) \cdot C^{-1} dY_s - \frac{1}{2} \int_0^t h(X_s) \cdot C^{-1} h(X_s) ds \right).$$

We can now introduce the unnormalised filtering measures

$$\rho_t[\phi] = \mathbb{E} [\phi(X_t) l_t | \mathcal{Y}_t], \quad t \geq 0.$$

The measures  $\rho_t$  satisfy the Zakai equation:

**Proposition A.1** (Zakai equation). *The evolution of  $(\rho_t)_{t \geq 0}$  is given by*

$$\rho_t[\phi] = \rho_0[\phi] + \int_0^t \rho_s[\mathcal{L}\phi] ds + \int_0^t \rho_s[\phi h] \cdot C^{-1} dY_s + \int_0^t \rho_s[\nabla \phi] \cdot B dY_s, \quad \mathbb{P} - a.s., \quad \forall \phi \in C_b^2(\mathbb{R}^D), \quad (103)$$

where

$$\mathcal{L}\phi = f \cdot \nabla \phi + \frac{1}{2} \text{Trace}(G \nabla^2 \phi)$$

is the generator associated to (1a). The filtering measures  $(\pi_t)_{t \geq 0}$  can be recovered from the Kallianpur-Striebel formula (or Bayes theorem)

$$\pi_t[\phi] = \frac{\rho_t[\phi]}{\rho_t[\mathbf{1}]} \quad (104)$$

*Proof.* This follows from the results in [10, Section 3.8].  $\square$

Next, we write the Zakai equation in its Stratonovich form:

**Lemma A.2.** *The Stratonovich-version of the Zakai equation is given by*

$$\begin{aligned} \rho_t[\phi] = & \rho_0[\phi] + \int_0^t \rho_s[\mathcal{L}\phi] ds + \int_0^t \rho_s[\phi h] C^{-1} \circ dY_s + \int_0^t \rho_s[\nabla \phi] \cdot B \circ dY_s \\ & - \frac{1}{2} \int_0^t \left( \rho_s[(\phi h)^T C^{-1} h] + \rho_s[\text{Trace}(\nabla(\phi h) \cdot B) + \nabla \phi \cdot B h] \right) ds \\ & - \frac{1}{2} \int_0^t \rho_s[\text{Trace}(D^2 \phi \underbrace{G^{1/2} U^T C^{-1} U G^{1/2}}_{BCB^T})] ds. \end{aligned}$$

*Proof.* Using (103), we see that for  $i = 1, \dots, d$

$$\rho_t[\phi h_i] = \sum_{j,k=1}^d \int_0^t \rho_s[\phi h_i h_j] (C^{-1})^{jk} dY_s^k + \sum_{r=1}^D \sum_{k=1}^d \int_0^t \rho_s[\partial_r(\phi h_i)] B^{rk} dY_s^k + FV,$$

where  $FV$  stands for a contribution of finite variation. Recalling  $\langle Y^k, Y^{\bar{k}} \rangle_t = C^{k\bar{k}} t$ , for  $k, \bar{k} = 1, \dots, d$ , we obtain

$$\int_0^t \rho_s[\phi h_i] C^{-1} dY_s = \int_0^t \rho_s[\phi h_i] C^{-1} \circ dY_s - \frac{1}{2} \int_0^t \sum_{i=1}^d \left( \sum_{j=1}^d \rho_s[\phi h_i h_j] (C^{-1})^{ij} + \sum_{r=1}^D \rho_s[\partial_j(\phi h_i) B^{ji}] \right) ds.$$

Similarly, for  $r = 1, \dots, D$ ,

$$\rho_t[\partial_r \phi] = \sum_{j,k=1}^d \int_0^t \rho_s[(\partial_r \phi) h_j] (C^{-1})^{jk} dY_s^k + \sum_{p=1}^D \sum_{k=1}^d \int_0^t \rho_s[\partial_r \partial_p \phi] B^{pk} dY_s^k + FV,$$

implying

$$\int_0^t \rho_s[\nabla \phi] B dY_s = \int_0^t \rho_s[\nabla \phi] B \circ dY_s - \frac{1}{2} \int_0^t \sum_{r=1}^D \left( \sum_{i=1}^d \rho_s[(\partial_r \phi) h_i] B^{ri} + \sum_{p=1}^D \rho_s[\partial_i \partial_p \phi] (BCB^T)^{rp} \right) ds.$$

□

We now proceed to the proof of Proposition 2.1:

*Proof of Proposition 2.1.* Let us define

$$\hat{\rho}_t[\phi] = \mathbb{E} \left[ \phi(\hat{X}_t) l_t | \mathcal{Y}_t \right], \quad t \geq 0,$$

which is the unnormalised filtering measure associated to (5). Since  $X_t$  and  $\hat{X}_t$  are independent given  $\mathcal{Y}_t$ , we have that

$$\hat{\rho}_t[\phi] = \mathbb{E} \left[ \phi(\hat{X}_t) | \mathcal{Y}_t \right] \mathbb{E} [l_t | \mathcal{Y}_t] = \hat{\pi}_t[\phi] \rho_t[\mathbf{1}].$$

From (104) it thus follows that  $\pi_t = \hat{\pi}_t$  is equivalent to  $\rho_t = \hat{\rho}_t$ . In the following we therefore compute the evolution of  $\hat{\rho}_t$ . Notice first that

$$M_t = \int_0^t h(X_s) \cdot C^{-1} dY_s = \int_0^t h(X_s) \cdot C^{-1} \circ dY_s - \frac{1}{2} \langle h(X) C^{-1}, Y \rangle_t,$$

where, using  $B = G^{\frac{1}{2}} U^T C^{-1}$ ,

$$\langle h(X) C^{-1}, Y \rangle_t = \sum_{i,j=1}^d \langle h^i(X) (C^{-1})_{ij}, Y^j \rangle_t = \sum_{r=1}^D \sum_{i=1}^d \int_0^t \partial_r h^i(X_s) B^{ri} ds = \int_0^t \text{Trace}(\nabla h(X_s) \cdot B) ds.$$

From  $dl_t = l_t dM_t$  we have

$$\begin{aligned} l_t &= 1 + \int_0^t l_s \circ dM_s - \frac{1}{2} \int_0^t l_s d\langle M \rangle_s \\ &= 1 + \int_0^t l_s h(X_s) \circ C^{-1} dY_s - \frac{1}{2} \int_0^t l_s \text{Trace}(\nabla h(X_s) \cdot B) ds - \frac{1}{2} \int_0^t l_s h(X_s) \cdot C^{-1} h(X_s) ds, \end{aligned}$$

implying

$$\phi(\hat{X}_t) l_t = \phi(\hat{X}_0) + \int_0^t \phi(\hat{X}_s) l_s h(X_s) \cdot C^{-1} \circ dY_s - \frac{1}{2} \int_0^t \phi(\hat{X}_s) l_s h(X_s) \cdot C^{-1} h(X_s) ds \quad (108a)$$

$$- \frac{1}{2} \int_0^t \phi(\hat{X}_s) l_s \text{Trace}(\nabla h(X_s) \cdot B) ds + \int_0^t l_s \nabla \phi(\hat{X}_s) \circ d\hat{X}_s. \quad (108b)$$

The last term satisfies

$$\begin{aligned} \int_0^t l_s \nabla \phi(\hat{X}_s) \circ d\hat{X}_s &= \int_0^t l_s \nabla \phi(\hat{X}_s) \cdot f(\hat{X}_s) ds + \int_0^t l_s \nabla \phi(\hat{X}_s) \circ G^{1/2} d\widehat{W}_s \\ &\quad + \int_0^t l_s \nabla \phi(\hat{X}_s) K_s(\hat{X}_s) C^{-1} \circ dY_s - \int_0^t l_s \nabla \phi(\hat{X}_s) \cdot K_s(\hat{X}_s) C^{-1} h(\hat{X}_s) ds \\ &\quad - \int_0^t l_s \nabla \phi(\hat{X}_s) \cdot K_s(\hat{X}_s) C^{-1} \circ (U d\widehat{W}_s + R^{1/2} d\widehat{V}_s) + \int_0^t l_s \nabla \phi(\hat{X}_s) \cdot \Xi_s(\hat{X}_s) ds. \end{aligned}$$

Next, we convert the terms involving  $\widehat{V}$  and  $\widehat{W}$  back to their Itô-form,

$$\begin{aligned} \int_0^t l_s \nabla \phi(\widehat{X}_s) \circ G^{1/2} d\widehat{W}_s &= \int_0^t l_s \nabla \phi(\widehat{X}_s) G^{1/2} d\widehat{W}_s \\ &+ \frac{1}{2} \int_0^t l_s \text{Trace}(D^2 \phi(\widehat{X}_s) G) ds - \frac{1}{2} \int_0^t l_s \text{Trace} \left( D^2 \phi(\widehat{X}_s) (K_s(\widehat{X}_s) B^T) \right) ds. \end{aligned}$$

Here we used the fact that, for every  $i, k = 1, \dots, D$  we have  $\langle l, (G^{1/2})^{ik} \widehat{W}^k \rangle_t = 0$  and

$$\langle \widehat{X}^j, (G^{1/2})^{ik} \widehat{W}^k \rangle_t = G^{ij} t - \int_0^t \left( K_s(\widehat{X}_s) C^{-1} U G^{1/2} \right)^{ji} ds,$$

where  $G^{1/2} U^T C^{-1} = B$ . Similarly,

$$\begin{aligned} \int_0^t l_s \nabla \phi(\widehat{X}_s) \cdot K_s(\widehat{X}_s) C^{-1} \circ \left( U d\widehat{W}_s + R^{1/2} d\widehat{V}_s \right) &= \int_0^t l_s \nabla \phi(\widehat{X}_s) \cdot K_s(\widehat{X}_s) C^{-1} \left( U d\widehat{W}_s + R^{1/2} d\widehat{V}_s \right) \\ &+ \frac{1}{2} \int_0^t l_s \text{Trace}[\nabla(\nabla \phi \cdot K_s)(\widehat{X}_s) \cdot C^{-1} (G^{1/2} U^T - K_s(\widehat{X}_s))] ds, \end{aligned}$$

using  $C = U U^T + R$  and, for  $k = 1, \dots, D$  and  $l = 1, \dots, d$ ,

$$\langle \widehat{X}^k, (U \widehat{W} + R^{1/2} \widehat{V})^l \rangle_t = (G^{1/2} U^T)^{kl} t - \int_0^t K_s^{kl}(\widehat{X}_s) ds.$$

We now take the conditional expectation in (108). Note that the conditional expectation w.r.t.  $\mathcal{Y}$  commutes with the integration in  $dY$  and that the conditional expectation of the integrals in  $dW$  and  $dV$  vanishes because  $(W, V)$  are independent from  $\mathcal{Y}$ , see [26, Appendix B]. We obtain

$$\widehat{\rho}_t[\phi] = \widehat{\rho}_0[\phi] + \int_0^t \widehat{\rho}_s[\phi] \pi_s[h] \cdot C^{-1} \circ dY_s - \frac{1}{2} \int_0^t \widehat{\rho}_s[\phi] \pi_s[h \cdot C^{-1} h] ds \quad (112a)$$

$$+ \int_0^t \widehat{\rho}_s[\mathcal{L}\phi] ds - \frac{1}{2} \int_0^t \widehat{\rho}_s[\text{Trace}(D^2 \phi K_s B^T)] ds + \int_0^t \widehat{\rho}_s[\nabla \phi \cdot K_s] C^{-1} \circ dY_s \quad (112b)$$

$$- \int_0^t \widehat{\rho}_s[\nabla \phi \cdot K_s C^{-1} h] ds - \frac{1}{2} \int_0^t \widehat{\rho}_s[\text{Trace}(\nabla(\nabla \phi \cdot K_s) \cdot C^{-1} (G^{1/2} U^T - K_s))] ds. \quad (112c)$$

$$- \frac{1}{2} \int_0^t \widehat{\rho}_s[\phi] \pi_s[\text{Trace}(DhB)] ds + \int_0^t \widehat{\rho}_s[\nabla \phi \cdot \Xi] ds \quad (112d)$$

Importantly, we have used the fact that  $X_t$  and  $\widehat{X}_t$  are independent given  $\mathcal{Y}_t$ , and also that  $\widehat{W}_t$  and  $\widehat{V}_t$  are independent from  $\mathcal{Y}_t$ . The next step is to compare (105) and (112). The  $dY$ -contributions agree if and only if

$$\rho_s[\phi h] + \rho_s[\nabla \phi] \cdot BC = \widehat{\rho}_s[\phi] \pi_s[h] + \widehat{\rho}_s[\nabla \phi \cdot K_s],$$

which we recognise (after identifying  $\widehat{\pi} = \pi$  and  $\widehat{\rho} = \rho$ ) to be a weak version of (6).

To compare the  $ds$ -contributions, let us first manipulate the second term in (112c) using (6),

$$\begin{aligned} \widehat{\rho}_s[\partial_m(\partial_i \phi K_s^{ij})(C^{-1})^{jk}(G^{1/2} U^T - K_s)^{mk}] &= - \int (\partial_i \phi K_s^{ij})(C^{-1})^{jk} \partial_m \left( \widehat{\rho}_s(G^{1/2} U^T - K_s)^{mk} \right) dx \\ &= - \int (\partial_i \phi K_s^{ij})(C^{-1})^{jk} \left( h^k - \widehat{\pi}_s[h^k] \right) d\widehat{\rho}_s = -\widehat{\rho}_s[\nabla \phi \cdot K_s C^{-1} (h - \widehat{\pi}_s[h])]. \end{aligned}$$

Similarly, the middle term in (112b) satisfies

$$\begin{aligned} \widehat{\rho}_s[(\partial_i \partial_j \phi) K_s^{il}] B^{jl} &= -B^{jl} \int (\partial_j \phi) \partial_i \left( \widehat{\rho}_s K_s^{il} \right) dx = B^{jl} \widehat{\rho}_s \left[ (\partial_j \phi) \left( h^l - \widehat{\pi}_s[h^l] \right) \right] - B^{jl} \int (\partial_j \phi) \partial_i \left( \widehat{\rho}_s (BC)^{il} \right) dx \\ &= B^{jl} \widehat{\rho}_s \left[ (\partial_j \phi) \left( h^l - \widehat{\pi}_s[h^l] \right) \right] + \widehat{\rho}_s[\partial_i \partial_j \phi] (BCB^T)^{ij} = \widehat{\rho}_s \left[ \nabla \phi \cdot B (h - \widehat{\pi}_s[h]) \right] + \widehat{\rho}_s[\text{Trace}(D^2 \phi (BCB^T))]. \end{aligned}$$

We now collect terms and compare the  $ds$ -contributions in (105) and (112), arriving at

$$- \frac{1}{2} \left( \rho_s[\phi h \cdot C^{-1} h] + \rho_s[\text{Trace}(D(\phi h)B) + \nabla \phi \cdot Bh] \right) \quad (115a)$$

$$= - \frac{1}{2} \widehat{\rho}_s[\phi] \pi_s[h \cdot C^{-1} h] - \frac{1}{2} \widehat{\rho}_s \left[ \nabla \phi \cdot B (h - \widehat{\pi}_s[h]) \right] \quad (115b)$$

$$- \frac{1}{2} \widehat{\rho}_s \left[ \nabla \phi \cdot K_s C^{-1} (h + \widehat{\pi}_s[h]) \right] + \widehat{\rho}_s[\nabla \phi \cdot \Xi] - \frac{1}{2} \widehat{\rho}_s[\phi] \pi_s[\text{Trace}(DhB)]. \quad (115c)$$

Next, we work on the first term in (115c),

$$\begin{aligned}
& \hat{\rho}_s \left[ (\partial_i \phi K_s^{ij})(C^{-1})^{jk} (h^k + \hat{\pi}_s[h^k]) \right] = - \int \phi \partial_i (\hat{\rho}_s K_s^{ij})(C^{-1})^{jk} (h^k + \hat{\pi}_s[h^k]) dx - \hat{\rho}_s \left[ \phi K_s^{ij}(C^{-1})^{jk} \partial_i h^k \right] \\
& = \int \phi (h^j - \pi[h^j])(C^{-1})^{jk} (h^k + \pi[h^k]) d\hat{\rho}_s - \int \phi \partial_i (\hat{\rho}_s (BC)^{ij})(C^{-1})^{jk} (h^k + \hat{\pi}_s[h^k]) dx - \hat{\rho}_s \left[ \phi K_s^{ij}(C^{-1})^{jk} \partial_i h^k \right] \\
& = \hat{\rho}_s [\phi (h^j - \pi[h^j])(C^{-1})^{jk} (h^k + \hat{\pi}_s[h^k])] + \hat{\rho}_s [\partial_i \phi (h^k + \hat{\pi}_s[h^k])] B^{jk} + \hat{\rho}_s [\phi B^{ik} \partial_i h^k] - \hat{\rho}_s \left[ \phi K_s^{ij}(C^{-1})^{jk} \partial_i h^k \right] \\
& = \hat{\rho}_s [\phi h^j h^k](C^{-1})^{jk} - \hat{\rho}_s [\phi] \pi_s[h^j] \pi_s[h^k](C^{-1})^{jk} + \hat{\rho}_s [\partial_i \phi (h^k + \hat{\pi}_s[h^k])] B^{jk} + \hat{\rho}_s [\phi B^{ik} \partial_i h^k] - \hat{\rho}_s \left[ \phi K_s^{ij}(C^{-1})^{jk} \partial_i h^k \right] \\
& = \hat{\rho}_s [\phi h C^{-1} h] - \hat{\rho}_s [\phi] \pi_s[h] C^{-1} \pi_s[h] + \hat{\rho}_s [\nabla \phi \cdot B(h + \hat{\pi}_s[h])] + \hat{\rho}_s [\phi \text{Trace}(DhB)] - \hat{\rho}_s \left[ \phi \text{Trace}(K_s C^{-1} Dh^T) \right]
\end{aligned}$$

Plugging this into (115) we see that after a great number of cancellations that (115) reduces to

$$\rho_s[\nabla \phi \cdot \Xi_s] = \frac{1}{2} \rho_s[\phi] \left( \pi[\text{Trace}(Dh \cdot B) - h C^{-1} h] - \pi[h] C^{-1} \pi[h] \right) - \frac{1}{2} \rho_s[\phi \text{Trace}(K_s C^{-1} Dh)].$$

Using (6) in its weak formulation tested against  $h$  on the first term in the right-hand side we obtain

$$\pi[\nabla \phi \cdot \Xi_s] = \frac{1}{2} \left( \pi[\phi] \pi[\text{Trace}(K C^{-1} Dh)] - \pi[\phi \text{Trace}(K C^{-1} Dh)] \right), \quad (117)$$

which is a weak version of (7).

We have thus obtained constraints on the coefficients of  $K$  and  $\Xi$  of the McKean-Vlasov equation (5) such that  $\hat{\rho}$  solves the Zakai equation satisfied by the conditional density  $\rho$  of the signal of the filtering problem. We can conclude because the solution to the Zakai equation is assumed to be unique.  $\square$

## B On the connection between McKean-Vlasov filtering and maximum likelihood estimation

Here we discuss the relationship between parameter estimation based on the ensemble Kalman filter and the maximum likelihood approach considered in [37]. In a nutshell, the two methods essentially agree in the noiseless case  $R = 0$  with Gaussian initial condition, as relevant McKean-Vlasov dynamics can be solved explicitly in this case. This connection has already been pointed out in [80].

We are given a  $d$ -dimensional process  $Z$ , which depends on an  $m$ -dimensional parameter  $\theta$ . We observe  $Z$  through the observation  $Y$ , which is possibly noisy.

$$dZ_t = g(Z_t) \theta dt + \tilde{G}^{\frac{1}{2}} dW_t, \quad (118a)$$

$$d\theta_t = 0, \quad (118b)$$

$$dY_t = dZ_t + R^{\frac{1}{2}} dV_t. \quad (118c)$$

Here  $W$  and  $V$  are independent  $d$ -dimensional Brownian Motions,  $g : \mathbb{R}^d \rightarrow \mathbb{R}^{d \times m}$  is a given  $C_b^2$  function and  $\tilde{G} \in \mathbb{R}^{d \times d}$  is a given positive semidefinite deterministic matrix.

Let  $D = d + m$ . We set the  $D$ -dimensional signal process as  $X := (Z, \theta)$  and replace the equation for  $dZ$  into the equation for the observation  $dY$ . System (118) becomes equivalent to system (1) with the following drift coefficients

$$f(x) = f(z, \theta) = (g(z) \theta, 0), \quad h(x) = h(z, \theta) = g(z) \theta.$$

Moreover, the coefficients of the noise are

$$G = \begin{pmatrix} \tilde{G} & 0 \\ 0 & 0 \end{pmatrix}, \quad U = \begin{pmatrix} \tilde{G}^{\frac{1}{2}} & 0 \end{pmatrix}, \quad G = \tilde{G}.$$

Remember  $\pi_t[\phi] := \mathbb{E}[\phi(X_t) | \mathcal{Y}_t]$  is the conditional law of  $X$  given  $Y$ .

First we consider noiseless observations, i.e. we set  $R = 0$ . In this case we have that

$$C := UU^\top + R = \tilde{G}, \quad B := G^{\frac{1}{2}} U^\top C^{-1} = \begin{pmatrix} \text{Id}_{d \times d} \\ 0_{m \times m} \end{pmatrix}.$$

To convert system (118) to the McKean-Vlasov equation we need to compute  $P$  as defined in (8), which can be decomposed as follows

$$P(\pi) = \begin{pmatrix} \text{Cov}_\pi(z, h) \\ \text{Cov}_\pi(\theta, h) \end{pmatrix} C^{-1} + B := \begin{pmatrix} \pi \left[ z(h - \pi[h])^\top \right] \\ \pi \left[ \theta(h - \pi[h])^\top \right] \end{pmatrix} C^{-1} + B.$$

In this case with  $R = 0$ , we have  $Y = Z$ , which implies

$$\pi_t(dz, d\theta) = \delta(z - Z_t)\pi_t^\theta(d\theta), \quad \pi_t[h] = \int_{\mathbb{R}^D} g(z)\theta\pi_t(dz, d\theta) = g(Z_t) \int_{\mathbb{R}^m} \theta\pi_t^\theta(d\theta) =: g(Z_t)\bar{\theta}_t$$

We have the following

$$\begin{aligned} \text{Cov}_{\pi_t}(z, h) &= \int_{\mathbb{R}^{d+m}} z (g(z)\theta - g(Z_t)\bar{\theta}_t)^\top \pi_t(dz, d\theta) = Z_t \left( \int_{\mathbb{R}^m} (\theta - \bar{\theta}_t)\pi_t^\theta(d\theta) \right)^\top g(Z_t)^\top = 0_{d \times d}. \\ \text{Cov}_{\pi_t}(\theta, h) &= \int_{\mathbb{R}^m} \theta (\theta - \bar{\theta}_t)^\top \pi_t^\theta(d\theta) g(Z_t)^\top = \text{Var}(\theta_t) g(Z_t)^\top. \end{aligned}$$

We have thus obtained  $P$  explicitly

$$P(\pi_t) = \begin{pmatrix} \text{Cov}_{\pi_t}(z, h) \\ \text{Cov}_{\pi_t}(\theta, h) \end{pmatrix} C^{-1} + B = \begin{pmatrix} 0_{d \times d} \\ \text{Var}(\theta_t)g(Z_t)^\top \end{pmatrix} \tilde{G}^{-1} + \begin{pmatrix} \text{Id}_{d \times d} \\ 0_{m \times m} \end{pmatrix} = \begin{pmatrix} \text{Id}_{d \times d} \\ \text{Var}(\theta_t)g(Z_t)^\top \tilde{G}^{-1} \end{pmatrix}$$

We now compute  $\hat{\Gamma}$ , which is defined in (9). We start with the  $z$  contribution, for  $\gamma = 1, \dots, d$ , we have

$$\pi[z^\gamma(Dh - \pi[Dh])]^\top = Z_t^\gamma \pi[Dh - \pi[Dh]]^\top = 0_{D \times d}.$$

Now compute the contribution in  $\theta$ . For  $\gamma = 1, \dots, m$  and  $i = 1, \dots, d$ , we have

$$\pi_t \left[ \bar{\theta}^\gamma Dh^\top \right]_{j,i} = \sum_{k=1}^m \partial_{z^j} g^{i,k}(Z_t) \text{Cov}(\theta_t^k, \theta_t^\gamma) 1_{j \leq d} = (\partial_{z^j} g(Z_t) \text{Var}(\theta_t))^{i,\gamma} 1_{j \leq d} \quad j = 1, \dots, d+m.$$

So that we have

$$P(\pi_t)^\top \pi_t \left[ \theta^\gamma (Dh - \pi_t[Dh])^\top \right] = \begin{pmatrix} \text{Id}_{d \times d} & (\text{Var}(\theta_t)g(Z_t)^\top \tilde{G}^{-1})^\top \\ 0_{m \times d} & \end{pmatrix} \begin{pmatrix} Dg(Z_t)^\top \text{Var}(\theta_t)^{-\gamma} \\ 0_{m \times d} \end{pmatrix} = Dg(Z_t)^\top \text{Var}(\theta_t)^{-\gamma}$$

and

$$\Gamma^\gamma(\pi_t) = -\frac{1}{2} \text{Trace} (Dg(Z_t) \text{Var}(\theta_t)^{-\gamma}).$$

If  $m = 1$ , we have

$$\Gamma(\pi_t) = \begin{pmatrix} 0_d \\ -\frac{1}{2} \text{Var}(\theta_t) \text{Trace} Dg(Z_t) \end{pmatrix}.$$

We can now write the McKean-Vlasov dynamics (11),

$$dZ_t = d\mathbf{Y}_t, \tag{119a}$$

$$d\hat{\theta}_t = \text{Var}(\hat{\theta}_t)g(Z_t)^\top \tilde{G}^{-1} \left[ d\mathbf{Y}_t - \tilde{G}^{\frac{1}{2}} d\widehat{W}_t - g(Z_t)\hat{\theta}_t dt \right] - \frac{1}{2} \text{Var}(\hat{\theta}_t) \text{Trace} Dg(Z_t) dt. \tag{119b}$$

We interpret equation (119) in the sense of Definition 4.12. Taking the expectation in (119b), and writing this equation in integral form, we obtain the rough path maximum likelihood estimator from [37], up to the rescaling  $\text{Var}(\hat{\theta}_t)$ , see also [80].

We would like to obtain an equation for the variance of  $\hat{\theta}$ . We start by computing the following equation for the centered random variable  $\bar{\theta} := \hat{\theta} - \pi[\theta]$ ,

$$d\bar{\theta}_t = -\text{Var}(\hat{\theta}_t)g(Z_t)^\top \tilde{G}^{-1} \left[ \tilde{G}^{\frac{1}{2}} d\widehat{W}_t + g(Z_t)\bar{\theta}_t dt \right].$$

By taking the square and using Itô formula we obtain the following equation for  $\text{Var}(\hat{\theta})$

$$d \text{Var}(\hat{\theta}_t) = -\text{Var}(\hat{\theta}_t)^2 g(\mathbf{Y}_t)^\top \tilde{G}^{-1} g(\mathbf{Y}_t) dt.$$

The solution is

$$\text{Var}(\hat{\theta}_t) = \left( \int_0^t g(\mathbf{Y}_s)^\top \tilde{G}^{-1} g(\mathbf{Y}_s) ds + \frac{1}{\text{Var}(\hat{\theta}_0)} \right)^{-1}.$$

Assume now that there exists  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  such that  $g = \tilde{G}^\top \nabla f$ . Since the variance is a path of bounded variation, we have that the integrand in the rough integral in equation (119b) is controlled by  $\mathbf{Y}$  in the following way

$$\begin{aligned} \text{Var}(\hat{\theta}_t)g(Y_t)^\top \tilde{G}^{-1} - \text{Var}(\hat{\theta}_s)g(Y_s)^\top \tilde{G}^{-1} &= \text{Var}(\hat{\theta}_t) \nabla f(Y_t)^\top - \text{Var}(\hat{\theta}_s) \nabla f(Y_s)^\top \\ &= \text{Var}(\hat{\theta}_t) D^2 f(Y_t) \delta Y_{s,t} + R_{s,t}. \end{aligned}$$

The Gubinelli derivative  $\text{Var}(\hat{\theta}_t) D^2 f(Z_t)$  is symmetric, which means that the rough integral in (119b) does not depend on the area of the geometric rough path  $\mathbf{Y}$ . Hence, the expansion of the integral is completely determined by the path itself.

## C Bibliography

- [1] A. Abdulle, G. Garegnani, G. A. Pavliotis, A. M. Stuart, and A. Zaroni. Drift estimation of multiscale diffusions based on filtered data. *arXiv preprint arXiv:2009.13457*, 2020.
- [2] Y. Ait-Sahalia, P. A. Mykland, and L. Zhang. How often to sample a continuous-time process in the presence of market microstructure noise. *The review of financial studies*, 18(2):351–416, 2005.
- [3] A. L. Allan. Robust filtering and propagation of uncertainty in hidden Markov models. *arXiv preprint arXiv:2005.04982*, 2020.
- [4] A. L. Allan, S. N. Cohen, et al. Pathwise stochastic control with applications to robust filtering. *Annals of Applied Probability*, 30(5):2274–2310, 2020.
- [5] R. Azencott, A. Beri, A. Jain, and I. Timofeyev. Sub-sampling and parametric estimation for multiscale dynamics. *Communications in Mathematical Sciences*, 11(4):939–970, 2013.
- [6] R. Azencott, A. Beri, and I. Timofeyev. Adaptive sub-sampling for parametric estimation of Gaussian diffusions. *Journal of Statistical Physics*, 139(6):1066–1089, 2010.
- [7] I. Bailleul, R. Catellier, and F. Delarue. Propagation of chaos for mean field rough differential equations. *arXiv preprint arXiv:1907.00578*, 2019.
- [8] I. Bailleul, R. Catellier, and F. Delarue. Solving mean field rough differential equations. *Electron. J. Probab.*, 25:Paper No. 21, 51, 2020.
- [9] I. Bailleul and J. Diehl. The inverse problem for rough controlled differential equations. *SIAM Journal on Control and Optimization*, 53(5):2762–2780, 2015.
- [10] A. Bain and D. Crisan. *Fundamentals of stochastic filtering*, volume 60. Springer Science & Business Media, 2008.
- [11] P. Bálint and I. Melbourne. Statistical properties for flows with unbounded roof function, including the Lorenz attractor. *Journal of Statistical Physics*, 172(4):1101–1126, 2018.
- [12] R. Beeson, N. Namachchivaya, and N. Perkowski. Approximation of the filter equation for multiple timescale, correlated, nonlinear systems. *arXiv preprint arXiv:2010.16401*, 2020.
- [13] R. Beeson, N. Namachchivaya, and N. Perkowski. Quantitative convergence of the filter solution for multiple timescale nonlinear systems with coarse-grain correlated noise. *arXiv preprint arXiv:2011.12801*, 2020.
- [14] R. Beeson, H. C. Yeong, N. S. Namachchivaya, and N. Perkowski. Reduced order nonlinear filters for multi-scale systems with correlated sensor noise. In *2018 21st International Conference on Information Fusion (FUSION)*, pages 131–141. IEEE, 2018.
- [15] T. Berry and T. Sauer. Correlation between system and observation errors in data assimilation. *Monthly Weather Review*, 146(9):2913–2931, 2018.
- [16] A. N. Bishop and P. Del Moral. On the mathematical theory of ensemble (linear-gaussian) Kalman-Bucy filtering. *arXiv preprint arXiv:2006.08843*, 2020.
- [17] G. Burgers, P. Jan van Leeuwen, and G. Evensen. Analysis scheme in the ensemble Kalman filter. *Monthly weather review*, 126(6):1719–1724, 1998.
- [18] R. Carmona and F. Delarue. *Probabilistic theory of mean field games with applications. II*, volume 84 of *Probability Theory and Stochastic Modelling*. Springer, Cham, 2018. Mean field games with common noise and master equations.
- [19] T. Cass and T. Lyons. Evolving communities with individual preferences. *Proc. Lond. Math. Soc. (3)*, 110(1):83–107, 2015.
- [20] I. Chevyrev, P. K. Friz, A. Korepanov, I. Melbourne, and H. Zhang. Multiscale systems, homogenization, and rough paths. In *International Conference in Honor of the 75th Birthday of SRS Varadhan*, pages 17–48. Springer, 2016.
- [21] I. Chevyrev and A. Kormilitzin. A primer on the signature method in machine learning. *arXiv preprint arXiv:1603.03788*, 2016.

- [22] J. Clark. The design of robust approximations to the stochastic differential equations of nonlinear filtering. *Communication systems and random process theory*, 25:721–734, 1978.
- [23] J. Clark and D. Crisan. On a robust version of the integral representation formula of nonlinear filtering. *Probability theory and related fields*, 133(1):43–56, 2005.
- [24] M. Coghi, J.-D. Deuschel, P. K. Friz, and M. Maurelli. Pathwise McKean-Vlasov theory with additive noise. *Ann. Appl. Probab.*, 30(5):2355–2392, 2020.
- [25] M. Coghi and F. Flandoli. Propagation of chaos for interacting particles subject to environmental noise. *Ann. Appl. Probab.*, 26(3):1407–1442, 2016.
- [26] M. Coghi and B. Gess. Stochastic nonlinear Fokker–Planck equations. *Nonlinear Analysis*, 187:259–278, 2019.
- [27] M. Coghi and T. Nilssen. Rough nonlocal diffusions. *arXiv preprint arXiv:1905.07270*, 2019.
- [28] C. Cotter, G. Pavliotis, et al. Estimating eddy diffusivities from noisy Lagrangian observations. *Communications in Mathematical Sciences*, 7(4):805–838, 2009.
- [29] D. Crisan, J. Diehl, P. K. Friz, H. Oberhauser, et al. Robust filtering: correlated noise and multidimensional observation. *The Annals of Applied Probability*, 23(5):2139–2160, 2013.
- [30] D. Crisan, A. Lobbe, and S. Ortiz-Latorre. Pathwise approximations for the solution of the non-linear filtering problem. *arXiv preprint arXiv:2101.03957*, 2021.
- [31] A. M. Davie. Differential equations driven by rough paths: an approach via discrete approximation. *Applied Mathematics Research eXpress*, 2008, 2008.
- [32] M. Davis. On a multiplicative functional transformation arising in nonlinear filtering theory. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, 54(2):125–139, 1980.
- [33] M. Davis and M. Spathopoulos. Pathwise nonlinear filtering for nondegenerate diffusions with noise correlation. *SIAM journal on control and optimization*, 25(2):260–278, 1987.
- [34] M. H. Davis. A pathwise solution of the equations of nonlinear filtering. *Theory of Probability & Its Applications*, 27(1):167–175, 1982.
- [35] M. H. Davis. Pathwise nonlinear filtering with correlated noise. *The Oxford Handbook of Nonlinear Filtering*, pages 403–424, 2011.
- [36] P. Del Moral. Feynman-kac formulae. In *Feynman-Kac Formulae*, pages 47–93. Springer, 2004.
- [37] J. Diehl, P. Friz, H. Mai, et al. Pathwise stability of likelihood estimators for diffusions via rough paths. *The Annals of Applied Probability*, 26(4):2169–2192, 2016.
- [38] J. Diehl, P. K. Friz, and W. Stannat. Stochastic partial differential equations: a rough paths view on weak solutions via Feynman-Kac. *Ann. Fac. Sci. Toulouse Math. (6)*, 26(4):911–947, 2017.
- [39] R. L. Dobrušin. Vlasov equations. *Funktsional. Anal. i Prilozhen.*, 13(2):48–58, 96, 1979.
- [40] A. Doucet and A. M. Johansen. A tutorial on particle filtering and smoothing: Fifteen years later. *Handbook of nonlinear filtering*, 12(656-704):3, 2009.
- [41] A. Duncan, N. Nüsken, and L. Szpruch. On the geometry of stein variational gradient descent. *arXiv preprint arXiv:1912.00894*, 2019.
- [42] G. Evensen. The ensemble Kalman filter: Theoretical formulation and practical implementation. *Ocean dynamics*, 53(4):343–367, 2003.
- [43] G. Evensen. *Data assimilation: the ensemble Kalman filter*. Springer Science & Business Media, 2009.
- [44] G. Flint, B. Hambly, and T. Lyons. Discretely sampled signals and the rough hoff process. *Stochastic Processes and their Applications*, 126(9):2593–2614, 2016.
- [45] P. Florchinger. Zakai equation of nonlinear filtering with unbounded coefficients. the case of dependent noises. *Systems & control letters*, 21(5):413–422, 1993.

- [46] N. Fournier and A. Guillin. On the rate of convergence in Wasserstein distance of the empirical measure. *Probability Theory and Related Fields*, 162(3):707–738, 2015.
- [47] P. Friz, P. Gassiat, and T. Lyons. Physical Brownian motion in a magnetic field as a rough path. *Transactions of the American Mathematical Society*, 367(11):7939–7955, 2015.
- [48] P. K. Friz and M. Hairer. *A course on rough paths*. Springer, 2020.
- [49] P. K. Friz, A. Hocquet, and K. Lê. Rough stochastic differential equations. *arXiv preprint arXiv:2106.10340*, 2021.
- [50] P. K. Friz and N. B. Victoir. *Multidimensional stochastic processes as rough paths: theory and applications*, volume 120. Cambridge University Press, 2010.
- [51] S. Gailus and K. Spiliopoulos. Statistical inference for perturbed multiscale dynamical systems. *Stochastic Processes and their Applications*, 127(2):419–448, 2017.
- [52] S. Gailus and K. Spiliopoulos. Discrete-time statistical inference for multiscale diffusions. *Multiscale Modeling & Simulation*, 16(4):1824–1858, 2018.
- [53] D. Givon, R. Kupferman, and A. Stuart. Extracting macroscopic dynamics: model problems and algorithms. *Nonlinearity*, 17(6):R55, 2004.
- [54] G. A. Gottwald and S. Reich. Supervised learning from noisy observations: Combining machine-learning techniques with data assimilation. *arXiv preprint arXiv:2007.07383*, 2020.
- [55] I. Gyöngy. On the approximation of stochastic partial differential equations ii. *Stochastics: An International Journal of Probability and Stochastic Processes*, 26(3):129–164, 1989.
- [56] W. R. Hammersley, D. Šiška, and Ł. Szpruch. Weak existence and uniqueness for McKean–Vlasov SDEs with common noise. *The Annals of Probability*, 49(2):527–555, 2021.
- [57] A. Hocquet and M. Hofmanová. An energy method for rough partial differential equations. *Journal of Differential Equations*, 265(4):1407–1466, 2018.
- [58] M. Holland and I. Melbourne. Central limit theorems and invariance principles for Lorenz attractors. *Journal of the London Mathematical Society*, 76(2):345–364, 2007.
- [59] P. Imkeller, N. S. Namachchivaya, N. Perkowski, and H. C. Yeong. A homogenization approach to multiscale filtering. *Procedia IUTAM*, 5:34–45, 2012.
- [60] P. Imkeller, N. S. Namachchivaya, N. Perkowski, H. C. Yeong, et al. Dimensional reduction in nonlinear filtering: a homogenization approach. *The Annals of Applied Probability*, 23(6):2290–2326, 2013.
- [61] S. Kalliadasis, S. Krumscheid, and G. A. Pavliotis. A new framework for extracting coarse-grained models from time series with multiscale structure. *Journal of Computational Physics*, 296:314–328, 2015.
- [62] R. Klein. Scale-dependent models for atmospheric flows. *Annual review of fluid mechanics*, 42:249–274, 2010.
- [63] S. Krumscheid, G. A. Pavliotis, and S. Kalliadasis. Semiparametric drift and diffusion estimation for multiscale diffusions. *Multiscale Modeling & Simulation*, 11(2):442–473, 2013.
- [64] S. Krumscheid, M. Pradas, G. Pavliotis, and S. Kalliadasis. Data-driven coarse graining in action: Modeling and prediction of complex systems. *Physical Review E*, 92(4):042139, 2015.
- [65] T. G. Kurtz and J. Xiong. Particle representations for a class of nonlinear SPDEs. *Stochastic Processes and their Applications*, 83(1):103–126, 1999.
- [66] T. G. Kurtz and J. Xiong. Numerical solutions for a class of SPDEs with application to filtering. In *Stochastics in finite and infinite dimensions*, Trends Math., pages 233–258. Birkhäuser Boston, Boston, MA, 2001.
- [67] H. J. Kushner. A robust discrete state approximation to the optimal nonlinear filter for a diffusion. *Stochastics*, 3(1-4):75–83, 1980.
- [68] Y. A. Kutoyants. *Statistical inference for ergodic diffusion processes*. Springer Science & Business Media, 2013.

- [69] A. Lejay and T. Lyons. On the importance of the Lévy area for studying the limits of functions of converging stochastic processes. application to homogenization. In *Current trends in potential theory*, volume 7. The Theta foundation/American Mathematical Society, 2003.
- [70] T. Lelièvre and G. Stoltz. Partial differential equations and stochastic methods in molecular dynamics. *Acta Numerica*, 25:681, 2016.
- [71] D. Levin, T. Lyons, and H. Ni. Learning from the past, predicting the statistics for the future, learning an evolving system. *arXiv preprint arXiv:1309.0260*, 2013.
- [72] N. Lingala, N. Perkowski, H. Yeong, N. S. Namachchivaya, and Z. Rapti. Optimal nudging in particle filters. *Probabilistic Engineering Mechanics*, 37:160–169, 2014.
- [73] N. Lingala, N. Sri Namachchivaya, N. Perkowski, and H. C. Yeong. Particle filtering in high-dimensional chaotic systems. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 22(4):047509, 2012.
- [74] R. S. Liptser and A. N. Shiryaev. *Statistics of random processes II: Applications*, volume 6. Springer Science & Business Media, 2013.
- [75] E. N. Lorenz. Deterministic nonperiodic flow. *Journal of atmospheric sciences*, 20(2):130–141, 1963.
- [76] T. Lyons. Rough paths, signatures and the modelling of functions on streams. *arXiv preprint arXiv:1405.4537*, 2014.
- [77] H. P. McKean, Jr. A class of Markov processes associated with nonlinear parabolic equations. *Proc. Nat. Acad. Sci. U.S.A.*, 56:1907–1911, 1966.
- [78] J. Nolen, G. A. Pavliotis, and A. M. Stuart. Multiscale modelling and inverse problems. In *Numerical analysis of multiscale problems*, pages 1–34. Springer, 2012.
- [79] N. Nüsken. Stein optimal transport for Bayesian inference.
- [80] N. Nüsken, S. Reich, and P. J. Rozdeba. State and parameter estimation from observed signal increments. *Entropy*, 21(5):505, 2019.
- [81] N. Nüsken and D. Renger. Stein variational gradient descent: many-particle and long-time asymptotics. *arXiv preprint arXiv:2102.12956*, 2021.
- [82] S. C. Olhede, A. M. Sykulski, and G. A. Pavliotis. Frequency domain estimation of integrated volatility for Itô processes in the presence of market-microstructure noise. *Multiscale Modeling & Simulation*, 8(2):393–427, 2010.
- [83] A. Papavasiliou, G. Pavliotis, and A. Stuart. Maximum likelihood drift estimation for multiscale diffusions. *Stochastic Processes and their Applications*, 119(10):3173–3210, 2009.
- [84] S. Pathiraja, S. Reich, and W. Stannat. McKean-Vlasov SDEs in nonlinear filtering. *arXiv preprint arXiv:2007.12658*, 2020.
- [85] S. Pathiraja and W. Stannat. Analysis of the feedback particle filter with diffusion map based approximation of the gain. 2021.
- [86] G. Pavliotis and A. Stuart. *Multiscale methods: averaging and homogenization*. Springer Science & Business Media, 2008.
- [87] G. A. Pavliotis. *Stochastic processes and applications: Diffusion Processes, the Fokker-Planck and Langevin Equations*, volume 60. Springer, 2014.
- [88] G. A. Pavliotis, Y. Pokern, and A. M. Stuart. Parameter estimation for multiscale diffusions: an overview. In *Statistical methods for stochastic differential equations*, volume 124 of *Monogr. Statist. Appl. Probab.*, pages 429–472. CRC Press, Boca Raton, FL, 2012.
- [89] G. A. Pavliotis and A. Stuart. Parameter estimation for multiscale diffusions. *Journal of Statistical Physics*, 127(4):741–781, 2007.
- [90] J. Pidstrigach and S. Reich. Affine-invariant ensemble transform methods for logistic regression. *arXiv preprint arXiv:2104.08061*, 2021.
- [91] S. Reich. Data assimilation: the Schrödinger perspective. *Acta Numerica*, 28:635–711, 2019.

- [92] S. Reich and C. Cotter. *Probabilistic forecasting and Bayesian data assimilation*. Cambridge University Press, 2015.
- [93] D. Simon. *Optimal state estimation: Kalman, H infinity, and nonlinear approaches*. John Wiley & Sons, 2006.
- [94] C. Sparrow. The Lorenz equations. *Edited by Arun V. Holden*, page 111, 1982.
- [95] A.-S. Sznitman. Topics in propagation of chaos. In *École d’Été de Probabilités de Saint-Flour XIX—1989*, volume 1464 of *Lecture Notes in Math.*, pages 165–251. Springer, Berlin, 1991.
- [96] A. Taghvaei, J. De Wiljes, P. G. Mehta, and S. Reich. Kalman filter and its modern extensions for the continuous-time nonlinear filtering problem. *Journal of Dynamic Systems, Measurement, and Control*, 140(3), 2018.
- [97] A. Taghvaei, P. G. Mehta, and S. P. Meyn. Diffusion map-based algorithm for gain function approximation in the feedback particle filter. *SIAM/ASA Journal on Uncertainty Quantification*, 8(3):1090–1117, 2020.
- [98] H. Tanaka. Limit theorems for certain diffusion processes with interaction. In *Stochastic analysis (Katata/Kyoto, 1982)*, volume 32 of *North-Holland Math. Library*, pages 469–488. North-Holland, Amsterdam, 1984.
- [99] J. Wouters and G. A. Gottwald. Stochastic model reduction for slow-fast systems with moderate time scale separation. *Multiscale Modeling & Simulation*, 17(4):1172–1188, 2019.
- [100] T. Yang, P. G. Mehta, and S. P. Meyn. Feedback particle filter. *IEEE transactions on Automatic control*, 58(10):2465–2480, 2013.
- [101] H. C. Yeong, R. Beeson, N. S. Namachchivaya, N. Perkowski, and P. W. Sauer. Dynamic data-driven adaptive observations in data assimilation for multi-scale systems. In *Handbook of Dynamic Data Driven Applications Systems*, pages 47–73. Springer, 2018.
- [102] H. C. Yeong, R. T. Beeson, N. S. Namachchivaya, and N. Perkowski. Particle filters with nudging in multiscale chaotic systems: with application to the Lorenz-96 atmospheric model. *Journal of Nonlinear Science*, pages 1–34, 2020.
- [103] Y. Ying, J. Maddison, and J. Vanneste. Bayesian inference of ocean diffusivity from Lagrangian trajectory data. *Ocean Modelling*, 140:101401, 2019.
- [104] L.-S. Young. What are SRB measures, and which dynamical systems have them? *Journal of statistical physics*, 108(5):733–754, 2002.
- [105] L. Zhang, P. A. Mykland, and Y. Aït-Sahalia. A tale of two time scales: Determining integrated volatility with noisy high-frequency data. *Journal of the American Statistical Association*, 100(472):1394–1411, 2005.